MDForLives

THE GLOBAL PULSE OF HEALTHCARE · SURGERY & PROCEDURAL CARE INSIGHT REPORT

Digital Postoperative Monitoring: What Surgeons Need Beyond More Data

Surgeon perspectives on remote follow-up, wearables, recovery apps, wound surveillance, alert ownership, workload, and what makes postoperative data actionable after discharge.

DISCHARGE → REMOTE SIGNALS → ALERT REVIEW → CLINICAL ACTION

147	6	72.1%	8976297
Survey records	Countries represented USA · UK · Canada · Italy · France · Germany	Completion rate	SGID

WHY THIS REPORT

Postoperative care is gaining more digital signals, but surgeons still need a reliable way to turn them into timely action

Postoperative deterioration often begins between scheduled clinical contacts. A wound changes at home, a patient becomes short of breath, hydration slips, mobility stalls, or pain no longer follows the expected recovery trajectory. Digital monitoring offers a way to make some of those changes visible earlier, especially as hospital stays shorten and more recovery happens outside the ward.

For surgeons, however, the central challenge is not simply collecting data. A remote signal has to be trustworthy enough to matter, reviewed by the right person, connected to a response protocol, and integrated with the clinical record without creating an unmanageable alert queue. The value of a wearable, app, photograph, or patient-reported outcome therefore depends on the service around it.

The MDForLives survey examines how surgeons say their use of postoperative digital monitoring has changed, which technologies are already in use, what they most want to detect earlier, how teams respond when a concerning signal appears, who reviews the data, what happens to workload, and which barriers still limit scale.

> **Survey frame:** 147 survey records with a 72.1% completion rate. Country metadata represents USA · UK · Canada · Italy · France · Germany. The survey export provides different response bases by question; multi-select questions are reported as selected in the source data. Administrative fields are excluded from the report.

CLINICAL CONTEXT

The evidence base is growing, but the clinical effect depends on the intervention, the operation, and the pathway around the technology

A 2024 JAMA Network Open systematic review and meta-analysis of perioperative telemedicine in abdominal surgery included 19 studies and 10,536 patients. Telemedicine interventions were associated with fewer readmissions and emergency-department visits, while complication rates were not significantly different from conventional care. The finding supports a practical distinction: digital follow-up can improve surveillance and access even when it does not prevent the complication itself.

More recent reviews describe similar opportunity with important limits. A 2025 systematic review of remote wound monitoring reported largely favorable findings across the studies it included, while a 2026 perioperative telehealth review concluded that telehealth has potential across the surgical journey but that patient selection and technological infrastructure still need optimization.

Connected devices also introduce operational and safety questions. The US FDA notes that wireless medical devices can enable remote access to patient data and real-time monitoring, while also requiring attention to safe use, cybersecurity, and reliable connectivity. These concerns align with the workflow, interoperability, data-security, and ownership issues raised by surgeons in the MDForLives survey.

External evidence provides context for the direction of travel; the MDForLives findings below describe what this respondent group reports about its own surgical practice, priorities, and implementation friction.

EXECUTIVE SUMMARY

Six signals that define the gap between collecting postoperative data and making it clinically useful

01

36.5%

Adoption is accelerating

Use of digital postoperative monitoring has increased significantly for more than one-third of respondents; another 35.7% report moderate growth.

02

65.7%

Wound complications are the leading early-detection target

Surgical-site or wound problems narrowly lead bleeding, cardiovascular, or respiratory deterioration as the complication surgeons most want to identify earlier.

03

57.4%

Virtual follow-up and patient-reported outcomes lead future value

Surgeons favor a model that keeps patient contact and recovery reporting inside the monitoring pathway, not only physiological sensing.

04

46.7%

A digital alert most often triggers patient contact or further assessment

The signal usually starts a clinical triage process rather than directly changing treatment.

05

54.2%

Evidence, cost, and reimbursement remain the biggest barrier

Integration, security, alert burden, staffing, adherence, and responsibility follow closely, showing that scale is a systems problem.

06

47.7%

Postoperative data mainly supports or confirms clinical assessment

Only 12.1% say it frequently leads to a management change, reinforcing the continuing role of surgical judgment.

01

ADOPTION CURVE

Digital postoperative monitoring is expanding, but practices are not moving at the same pace

36.5% report a significant increase in use over the past three years; another 35.7% report a moderate increase.

CONTEXT

Postoperative follow-up used to be organized mainly around discharge instructions, scheduled clinic visits, telephone contact, and patient-initiated escalation. Connected monitoring tools add another layer: structured symptoms, activity, wound images, vital signs, and virtual contact can now be collected between those traditional checkpoints.

WHAT THE DATA SHOWS

Use has increased significantly for 36.5% and moderately for 35.7%, while 27.8% say it has remained largely unchanged. The distribution points to broad momentum, but not to a single level of maturity across surgical services.

THE INSIGHT

Growth in use does not tell us whether monitoring is integrated, staffed, or linked to escalation. Two teams can both report greater digital use while one has a mature virtual ward and the other has a stand-alone app that clinicians review inconsistently.

WHY IT MATTERS

Implementation decisions therefore need to separate technology adoption from pathway adoption. The stronger benchmark is whether a service knows which patients are monitored, which signals matter, who reviews them, and what happens when a threshold is crossed.



02

CURRENT TECHNOLOGY MIX

The current toolset is broad, with recovery apps and wearable monitoring ahead of digital wound surveillance

50.0% currently use digital recovery apps, symptom tracking, or virtual follow-up; 45.5% use RPM platforms or wearable physiological monitoring.

CONTEXT

Different postoperative problems produce different data. A patient recovering after a joint procedure may generate mobility or activity information; a patient after abdominal surgery may report symptoms and wound change; a higher-risk patient may require physiological monitoring. That diversity makes it unlikely that one device will cover the whole recovery pathway.

WHAT THE DATA SHOWS

Digital recovery apps, symptom tracking, or virtual follow-up are selected by 50.0%. RPM platforms or wearable physiological monitoring follow at 45.5%. AI/predictive analytics or EHR-integrated surveillance are at 34.5%, and digital wound monitoring or imaging at 23.6%. Respondents could select up to three technologies.

THE INSIGHT

The current mix suggests that surgical digital monitoring is already combining communication and measurement. Recovery apps create a structured channel for symptoms and follow-up, while wearables extend observation into the home. Wound imaging remains less common despite wound complications being the leading early-detection priority later in the survey.

WHY IT MATTERS

That gap may matter for service design. If the most valued clinical target is a wound complication, but wound imaging is one of the less frequently used current tools, the pathway may have an opportunity to connect patient photographs or structured wound assessment more directly to surgical review.



03

FUTURE MONITORING MODEL

Surgeons want the next monitoring layer to integrate recovery data rather than simply add more sensors

57.4% choose virtual follow-up and patient-reported outcome tracking; 51.9% choose integrated platforms that combine multiple monitoring technologies.

CONTEXT

A physiologic threshold can identify risk, but recovery is often multidimensional. Pain, oral intake, bowel or bladder function, wound appearance, activity, fatigue, and the patient's own sense that something is wrong can all shape postoperative assessment. Digital systems become more useful when they can bring those signals into one view.

WHAT THE DATA SHOWS

Virtual follow-up and patient-reported outcome tracking lead at 57.4%. Integrated multi-technology platforms follow at 51.9%, AI/predictive analytics and digital wound assessment at 46.3%, and RPM with continuous physiological monitoring at 41.7%.

THE INSIGHT

The ranking places communication and integration above stand-alone continuous monitoring. Surgeons appear to be asking for context, not only measurement: a way to know what the patient is experiencing, combine it with objective data, and decide what requires attention.

WHY IT MATTERS

A future postoperative platform may therefore need to work like a clinical workspace rather than a device dashboard. The goal is to reduce fragmentation by aligning symptoms, patient-reported outcomes, images, vital signs, and prior clinical context in the same review process.



04

EARLY DETECTION TARGETS

The highest-value signals are the ones that could narrow the time to recognition of clinically important deterioration

65.7% most value earlier identification of surgical-site or wound complications; 63.9% select bleeding, cardiovascular, or respiratory deterioration.

CONTEXT

Postoperative monitoring has a finite attention budget. A system that alerts on every deviation may generate more noise than value, while a system that misses the complications surgeons most worry about does not earn trust. The clinical target therefore has to be explicit before the technology is chosen.

WHAT THE DATA SHOWS

Surgical-site or wound complications lead at 65.7%, followed closely by bleeding, cardiovascular, or respiratory deterioration at 63.9%. Pain, dehydration, or other physiological deterioration are selected by 43.5%, and delayed recovery or multiple complications through integrated monitoring by 38.9%. Respondents could select up to three.

THE INSIGHT

The top two categories share one feature: delay matters. A wound infection, bleeding event, respiratory decline, or cardiovascular deterioration can begin outside a scheduled review window. The perceived value of digital monitoring is strongest when it shortens that blind interval.

WHY IT MATTERS

This prioritization can guide program design. Rather than monitoring every available variable, a surgical service can define the complications it most needs to detect earlier, identify which combination of patient report, image, or physiological data best supports that goal, and build the escalation process around those signals.



05

FIRST ACTION AFTER ALERT

Most digital alerts begin a triage conversation rather than an automated treatment decision

46.7% most commonly contact the patient or arrange further assessment after a potentially concerning postoperative change is detected.

CONTEXT

An abnormal value or patient-reported symptom is only one part of a postoperative decision. Surgeons still need to understand timing, severity, associated symptoms, procedure-specific risk, baseline recovery, and whether the change is transient or progressive.

WHAT THE DATA SHOWS

Contacting the patient or arranging further assessment leads at 46.7%. Another 33.6% most commonly arrange urgent in-person review, testing, or imaging. Reviewing trends before acting accounts for 11.2%, while direct treatment adjustment or escalation is selected by 8.4%.

THE INSIGHT

Digital data is acting primarily as a trigger for assessment. That is clinically important because it positions remote monitoring as an earlier entry point into the decision pathway, not as a replacement for examination or diagnostic work-up.

WHY IT MATTERS

The best-designed alert therefore needs a response protocol that is as clear as the threshold itself. The reviewer should know which questions to ask, whether a virtual assessment is adequate, when imaging or laboratory testing is required, and when the patient needs urgent face-to-face care.



06

ALERT OWNERSHIP

Monitoring responsibility is split almost evenly between surgical teams and dedicated clinical reviewers

34.6% say the operating surgeon or surgical team primarily reviews alerts; 33.6% point to nursing, APP, or dedicated RPM/virtual-care teams.

CONTEXT

Remote monitoring extends the clinical day. A signal may arrive when the surgeon is operating, in clinic, off service, or responsible for multiple postoperative patients at once. The practical question is not only who has the expertise to interpret the signal, but who is reliably available to see it in time.

WHAT THE DATA SHOWS

Operating surgeons or surgical teams lead at 34.6%, closely followed by nursing, APP, or dedicated RPM/virtual-care teams at 33.6%. Shared multidisciplinary teams account for 18.7%, while 13.1% report no formal monitoring process or a process that varies by patient or procedure.

THE INSIGHT

There is no dominant ownership model. That may be appropriate because different procedures and risk levels require different pathways, but the lack of a clear standard also creates a safety question: a digital alert is only useful if someone is explicitly accountable for reviewing and routing it.

WHY IT MATTERS

Services can reduce ambiguity by defining review windows, backup coverage, escalation criteria, and the handoff between virtual-care staff and the operating team. Ownership should be designed before monitoring volume increases, not after alerts start accumulating.



07

WORKLOAD REALITY

Digital monitoring does not consistently reduce workload; for some teams it creates another layer of work

29.0% report no meaningful workload change, 26.2% report increased workload, and 23.4% report reduced workload.

CONTEXT

One promise of remote monitoring is that earlier information and virtual follow-up could replace unnecessary calls or visits. But every digital channel also creates new tasks: onboarding patients, troubleshooting technology, reviewing dashboards, responding to alerts, documenting decisions, and coordinating escalation.

WHAT THE DATA SHOWS

No meaningful change is the largest response at 29.0%. Increased workload follows at 26.2%, reduced workload at 23.4%, and 21.5% say it is too early to determine or they are not currently using digital monitoring.

THE INSIGHT

The workload distribution is almost evenly split rather than clearly favorable. That suggests efficiency is not an automatic property of the technology. It is an outcome of pathway design, automation quality, staffing, alert specificity, and whether digital interactions replace or simply add to existing work.

WHY IT MATTERS

A monitoring program should therefore measure staff time as deliberately as it measures patient engagement. If the system increases review burden without preventing low-value contacts or focusing attention on higher-risk patients, adoption may stall even when clinicians believe the data has potential.



08

BARRIERS TO SCALE

The scaling barrier is a stack of system constraints rather than one missing feature

54.2% cite limited evidence, cost, or reimbursement; 51.4% cite workflow integration, interoperability, or data-security issues.

CONTEXT

Pilot programs can survive on motivated clinicians and manual workarounds. Routine services cannot. At scale, remote monitoring has to compete for budget, integrate with existing clinical systems, create manageable alert volumes, fit available staffing, and work for patients with different levels of digital access and confidence.

WHAT THE DATA SHOWS

Limited evidence, cost, or reimbursement is selected by 54.2%. Workflow integration, interoperability, or data-security issues follow at 51.4%; alert burden, data overload, or lack of dedicated staff at 48.6%; and patient adherence, digital literacy, or unclear monitoring responsibility at 43.0%. Respondents could select up to three.

THE INSIGHT

All four barriers sit above 40%, which means the implementation problem is multidimensional. Solving one issue in isolation may not unlock scale if another part of the pathway remains weak. A platform can be clinically useful but still fail because the economics, staffing model, or data integration is unsustainable.

WHY IT MATTERS

This is where procurement and clinical design need to meet. Technology selection should include evidence quality, reimbursement or funding, EHR integration, cybersecurity, staffing, alert governance, patient usability, and defined clinical ownership - not only device accuracy or feature count.



09

DATA TO DECISION

Technology-generated data is most often a clinical support layer, not the final decision-maker

47.7% say postoperative technology data mainly supports or confirms clinical assessment; only 12.1% say it frequently leads to a change in management.

CONTEXT

Surgeons are used to combining multiple incomplete signals: symptoms, examination, procedure-specific risk, laboratory results, imaging, wound appearance, and the expected recovery trajectory. Digital postoperative data enters that same reasoning process rather than replacing it.

WHAT THE DATA SHOWS

Nearly half, 47.7%, say digital data mainly supports or confirms clinical assessment. Another 30.8% say it occasionally influences management. Frequent management change is reported by 12.1%, while 9.3% say the data rarely provides meaningful clinical value.

THE INSIGHT

The result helps define a realistic role for postoperative technology. Its value may be greatest in prompting earlier attention, confirming or challenging a clinical impression, and making recovery trends visible between visits. That is different from expecting the system to routinely make the decision itself.

WHY IT MATTERS

This distinction can improve both expectations and evaluation. A monitoring pathway should not be judged only by the number of treatment changes it produces. It can also create value by identifying who needs review sooner, supporting reassurance when recovery is on track, and giving the surgeon a more complete picture before the next decision.



OPEN RESPONSE LENS

When surgeons describe the monitoring improvement they want most, the recurring theme is continuous visibility with a human response behind it

The open-ended question asked surgeons to name one technology, capability, or improvement in digital monitoring that could make the greatest difference to patient recovery or earlier detection of complications. Responses were multilingual and varied in detail. The themes below summarize the substantive comments without creating percentages or treating brief comments as coded quantitative data.

Continuous vital-sign monitoring after discharge

Several responses described home monitoring of heart rate, blood pressure, oxygen saturation, temperature, respiratory rate, or activity. The desired capability was not a single spot check, but a continuous or repeated view of recovery outside the hospital.

Automated alerts and predictive detection

Respondents described AI-supported or automated alerts that could identify abnormal change early and direct the team toward possible infection, bleeding, respiratory, cardiovascular, or other deterioration.

Wound images as an early clinical signal

Postoperative wound photographs and earlier availability of wound images appeared repeatedly, including comments focused specifically on early detection of wound complications.

Connected post-discharge follow-up

Teleconsultation, remote follow-up after discharge, app-linked monitoring, and devices that remain connected to the care team were recurring practical requests.

Recovery needs more than vital signs

A smaller set of comments pointed to physiotherapy monitoring, activity, urine output, and postoperative laboratory information, suggesting interest in broader recovery signals beyond standard wearable data.

Actionability matters more than data volume

The more detailed comments consistently connected monitoring to a clinical response: detect deviation, alert the team, contact the patient, and intervene earlier. The desired outcome was not simply a richer dashboard but a shorter path from change to assessment.

PEOPLE ALSO ASK · TOPIC FAQ

Frequently asked questions about digital postoperative monitoring and remote surgical follow-up.

What is digital postoperative monitoring?

Digital postoperative monitoring uses connected tools such as recovery apps, remote patient monitoring platforms, wearables, symptom questionnaires, wound images, and virtual follow-up to track recovery after surgery. The aim is to identify meaningful changes between scheduled visits and connect those signals to an appropriate clinical response.

How does remote patient monitoring work after surgery?

Patients may submit symptoms, patient-reported outcomes, photographs, activity data, or physiological measurements from home. A surgical, nursing, virtual-care, or multidisciplinary team reviews the information using predefined workflows and decides whether the patient needs reassurance, additional assessment, testing, or escalation.

What postoperative complications can remote monitoring help identify?

Depending on the technology and pathway, remote monitoring may help surface wound concerns, abnormal vital signs, bleeding or cardiopulmonary deterioration, pain, dehydration, reduced mobility, or delayed recovery. A digital signal is not a diagnosis and should be interpreted alongside the clinical context.

Can digital monitoring replace in-person postoperative follow-up?

Not universally. Digital monitoring can extend observation beyond the hospital and support triage, but some symptoms, examinations, wound concerns, or abnormal trends still require direct clinical assessment, testing, imaging, or urgent in-person review.

What is digital wound monitoring?

Digital wound monitoring uses photographs, structured symptom reporting, or connected wound-assessment tools to follow healing after surgery. It can help clinicians review change over time, but image quality, patient technique, clinical thresholds, and escalation pathways remain important.

What limits wider use of remote monitoring after surgery?

Common implementation challenges include evidence and reimbursement, workflow integration, interoperability, data security, alert burden, staffing, patient adherence, digital literacy, and uncertainty about who owns the monitoring response. These issues can matter as much as the device itself.

SURGICAL MONITORING LENS

What this means at four points in a digital postoperative monitoring pathway.

The survey suggests that the most useful pathway starts by defining the clinical problem, then builds the technology, ownership, and escalation process around it. Four design choices stand out.

1. Define the complication before the device

Start with the postoperative problem the service most needs to detect earlier - wound change, cardiopulmonary deterioration, dehydration, pain, or delayed recovery - and choose the data source that best supports that use case.

2. Give every signal an owner

Specify who reviews incoming data, how often, what backup coverage exists, and when nursing, virtual-care, or surgical teams hand a patient to one another.

3. Pair thresholds with escalation rules

An alert should point to a next step: patient contact, virtual assessment, urgent in-person review, testing, imaging, or escalation. The threshold is only half of the clinical design.

4. Make the home pathway easy to use

Connected monitoring must work for real patients after discharge. Minimize device burden, support digital literacy, integrate data into clinician workflow, and ensure that patient-reported symptoms can sit beside objective measurements.

CLOSING PERSPECTIVE

Digital postoperative monitoring becomes clinically useful when the signal, the reviewer, and the next action are designed as one pathway

The survey captures a surgical field that is already moving beyond episodic follow-up. Most respondents report at least some increase in digital postoperative monitoring, and current practice spans recovery apps, wearables, virtual follow-up, AI-linked surveillance, and wound technologies. Surgeons see particular value in earlier visibility of wound complications and physiological deterioration after discharge.

Yet the findings also show why more technology does not automatically produce a safer or more efficient pathway. Alerts usually lead to patient contact or further assessment rather than direct treatment change. Responsibility is almost evenly split between surgical teams and nursing or virtual-care teams. Workload can fall, stay unchanged, or increase. And the leading barriers - evidence, economics, integration, security, alert burden, staffing, adherence, and ownership - are all system issues.

The clinical role of the data is equally important. Nearly half say technology-generated information mainly supports or confirms assessment, and only a minority say it frequently changes management. That is not necessarily a weakness. In surgery, a monitoring tool can add value by shortening the time to awareness, helping the team triage risk, and making the next conversation better informed.

The direction suggested by the data is therefore selective and integrated: monitor the complications where earlier recognition matters, combine objective signals with patient-reported recovery, route alerts to a named reviewer, define escalation before launch, and evaluate workload as seriously as clinical performance. The goal is not a postoperative dashboard with more numbers. It is a recovery pathway with fewer blind spots.

Prioritize time-sensitive signals
Focus monitoring on postoperative changes where earlier recognition can alter triage, assessment, or escalation.

Make clinical ownership visible
Define who reviews alerts, when they are reviewed, and how responsibility transfers across teams and time periods.

Measure the work created by monitoring
Track alert volume, review time, duplication, and whether digital contacts replace or simply add to existing follow-up.

Build one view of recovery
Bring symptoms, patient-reported outcomes, wound information, vital signs, and clinical history together so the reviewer sees context, not isolated data points.

Help future surgical digital-health insights reflect the realities of postoperative care.

Join MDForLives to share clinical perspectives on remote monitoring, postoperative recovery, virtual follow-up, digital wound care, workflow, and early detection.

Join the Panel

REFERENCES AND DATA SOURCE

1. MDForLives internal survey report. All Markets Report for SGID 8976297, Project: Engagement Campaign | Surgeons Insight Survey-Sep | MDForLives | GID: 9000 | Team: Internal.
2. Grygorian A, et al. Digital Health Interventions and Patient Safety in Abdominal Surgery: A Systematic Review and Meta-Analysis. JAMA Network Open. 2024.
3. Tanveer F, et al. Effectiveness of remote patient monitoring for wound care: A systematic review. Journal of the Pakistan Medical Association. 2025.
4. Yilmaz Celik A, et al. Telehealth Applications Used in Perioperative Patient Care: A Systematic Review. Journal of PeriAnesthesia Nursing. 2026.
5. US Food and Drug Administration. Wireless Medical Devices. Digital Health Center of Excellence.
6. Khan MM, Shah N. AI-driven wearable sensors for postoperative monitoring in surgical patients: A systematic review. Computers in Biology and Medicine. 2025.

External sources provide clinical context. Survey percentages represent MDForLives respondent selections and are reported directly from the supplied source export. Multi-select question percentages can total more than 100%.

MDForLives

MDForLives is a global healthcare insights platform where healthcare professionals, patients, and caregivers discover, share, and shape the future of healthcare. This report presents aggregated professional perspectives for educational and insight purposes.

The Global Pulse of Healthcare
© MDForLives