Cancer Survivorship Care: Balancing Recurrence Risk and Long-Term Toxicity

SILVIA PARADISO, MD · ALESSIA LANZA · PHYSICIST · LIANA TSIKLAURI · CORRESPONDENCE CARE

113 Oncologists surveyed

6 Countries represented USA · UK · Canada · Italy · France · Germany

73.5% Completion rate

9007321 Study ID

WHY THIS REPORT

Cancer follow-up changes after treatment, but the clinical work does not become quieter.

Survivorship care uses oncology teams to manage key timelines at once: the possibility of recurrence and the long-term consequences of treatment. A patient may have stable scans but persistent fatigue; another may have no evidence of disease but ongoing anxiety. Clinicians must balance surveillance with quality of life, using tools that can miss recurrence, overtest low-risk patients, or overlook treatment effects.

This creates a practical tension. Follow-up must be intensive enough to detect clinically important changes, but targeted enough to avoid unnecessary imaging, biopsies, and distress. Clinicians must also decide when symptoms should be investigated and when they reflect cancer survivorship rather than disease recurrence. Clear criteria for escalation and shared decision-making can help set realistic expectations.

The MDForum survey explores where oncology HCPs intentionally follow up, what concerns them most, which late effects most change management, how patients understand survivorship care efforts, and which emerging tools clinicians believe could improve the next phase of follow-up.

METHODOLOGY

A cross-market survey of professional oncology HCPs.

The survey included professional HCP respondents across the USA, UK, Canada, Italy, France, and Germany. The sources for estimated CLV survey results with the 113 completed responses. Quantitative questions were closed-ended, scored as respondents progressed through the survey.

The quantitative findings are reported separately as provided in the survey response export. The open-ended case question is analyzed qualitatively, with responses grouped into themes and illustrated with representative quotes.

Administrative tools such as free-charge tracking, fast-ID response, fast-ID, and incomplete statistics were excluded from the clinical analysis. Country-pooling was used only for geographically balanced.

EXECUTIVE SUMMARY

Six signals define the survivorship-care challenge.

01

46.1%

Higher-risk disease intensifies follow-up

Stage III–IV disease is the most common patient subgroup driving more frequent surveillance.

02

46.1%

Recurrence risk anchors surveillance intensity

It is the most common factor driving imaging-first clinic scenarios and surveillance.

03

44.3%

Persistent toxicity can change treatment

Dose reduction or modification is the leading reported response to ongoing treatment toxicity.

04

46.6%

Cardiotoxicity anchors survivorship strategy

It outranks cognitive impairment, neuropathy, and secondary malignancies as the most clinically important late effect.

05

54.0%

Confirmation is the biggest screen barrier

Fragmented oncology primary care coordination is the strongest operational signal in the survey.

06

32.2%

Recruitment awareness direction leads future priorities

AI-based recruitment risk prediction follows at 26.7%.

SURVIVORSHIP CARE OVERVIEW

Stage III–IV disease is the clearest signal for intensified surveillance follow-up.

46.1% clearly stage III–IV disease as the summer profile most warranting intensified follow-up

CONTEXT

Follow-up intensity cannot be uniform across all survivors. Disease burden, treatment exposure, and recurrence risk should shape how often patients are seen and how the follow-up plan is adjusted.

WHAT THIS SHOWS

Stage III–IV disease is the most common patient subgroup driving more frequent surveillance, with 46.1% clinicians selected it as the most common factor driving imaging-first clinic scenarios and surveillance.

THE SIGNAL

Respondents appear to give disease stage the greatest weight when deciding who needs intensified follow-up, while treatment intensity and unresolved toxicity are additional factors, not the primary trigger.

WHY IT MATTERS

Clear criteria help oncologists balance risk-based ongoing care decisions. Guidelines that only stage radiation toward disease progression may miss patients who have ongoing symptoms.

Survivor profile that most warrants intensified follow-up

Stage III–IV disease	46.1%
Treatment intensity	26.5%
Persistent toxicity	17.7%
Non-concerning symptoms	9.7%

Finding that most influences surveillance intensity

Recurrence risk	46.1%
Treatment toxicity	26.5%
Persistent toxicity	17.7%
New concerning symptoms	9.7%

RISK SURVEILLANCE & RECURRENCE

Recurrence risk is the central surveillance signal, ahead of stage, toxicity, and symptoms.

46.1% clearly recurrence risk as the finding that most influences surveillance intensity

CONTEXT

Post-treatment surveillance is meant to detect clinically meaningful recurrence while avoiding unnecessary testing. The difficult part is deciding which signal should drive imaging, labs, or clinic visits.

WHAT THIS SHOWS

Recurrence risk was the top survivorship finding, with 46.1% selecting it as the most influential factor in surveillance intensity. Treatment toxicity followed at 26.5%, and persistent toxicity at 17.7%.

THE SIGNAL

Respondents suggest that treatment-related toxicity is important, but the unresolved possibility of recurrence matters more than any single symptom or toxicity when setting the baseline surveillance pathway.

WHY IT MATTERS

Clear recurrence risk frameworks can support more consistent follow-up intensity while leaving room to escalate when new symptoms, toxicities, or clinician specific reasons emerge.

Finding that most influences surveillance intensity

Recurrence risk	46.1%
Treatment toxicity	26.5%
Persistent toxicity	17.7%
New concerning symptoms	9.7%

RISK SURVEILLANCE & RECURRENCE

Guideline-based imaging and individualized risk-based surveillance are almost evenly matched.

43.4% primarily use guideline-based imaging, while 39.8% use individualized risk-based surveillance.

CONTEXT

Surveillance creates consistency, but cancer survivorship rarely fits a single template. Stage, treatment history, recurrence biology, age, comorbidity, patient preference, and late effects may all justify adaptation.

WHAT THIS SHOWS

Guideline-based imaging leads at 43.4%, with individualized risk-based surveillance following at 39.8%. Symptom-driven surveillance is 16.8%, and symptom-driven follow-up is 9.7%.

THE SIGNAL

The small gap between the two leading approaches suggests that clinicians are not choosing between rigid schedules and fully personalized philosophies. They are complementary layers of the same follow-up process.

WHY IT MATTERS

Survivorship programs need enough structure to be reliable and enough flexibility to adapt to changes in recurrence risk, treatment toxicity, and patient context.

Primary surveillance strategy

Guideline-based imaging	43.4%
Individualized risk-based surveillance	39.8%
Symptom-driven follow-up	16.8%

RISK SURVEILLANCE & RECURRENCE

Uncertain recurrence risk leads a tightly clustered set of management constraints.

39.8% select uncertain recurrence risk, not capacity, comorbidity, and toxicity follow closely

CONTEXT

Survivorship management can be constrained by uncertainty even when the clinical picture looks stable. Of the 113 HCPs surveyed, 39.8% selected risk as the top limiter, immediately ahead of capacity, the operational constraint most often cited by clinicians.

WHAT THIS SHOWS

Uncertain recurrence risk is selected by 39.8%, limited follow-up capacity by 23.0%, comorbidity-driven by 19.5%, and late-treatment toxicity by 17.7%.

THE SIGNAL

No single issue dominates. The most even distribution suggests that survivorship decisions are rarely driven by one factor. Clinicians weigh risk and operational constraints rather than one isolated barrier point.

WHY IT MATTERS

Improving survivorship care therefore requires both better risk stratification and protected capacity to act on that risk.

What most limits effective survivorship management

Uncertain recurrence risk	39.8%
Limited follow-up capacity	23.0%
Comorbidity-driven toxicity	19.5%
Late-treatment toxicity	17.7%

RISK SURVEILLANCE & RECURRENCE

Dose reduction or modification is the most selected response to ongoing treatment toxicity.

44.3% choose dose reduction or modification, compared with 23.0% selecting symptom-directed pharmacotherapy.

CONTEXT

Persistent toxicity can blur the boundary between active treatment and survivorship. The clinical team may be trying to preserve treatment benefit while reducing the burden of long-term toxicity, quality of life, or long-term organ health.

WHAT THIS SHOWS

Dose reduction or modification leads at 44.3%, followed by supportive care at 23.0%, symptom-directed pharmacotherapy at 19.5%, and treatment withdrawal at 13.3%.

THE SIGNAL

Cardiotoxicity is actively another symptom burden. It can create a possible chronic disease pathway that changes how the survivor is monitored and which clinicians need to be involved.

WHY IT MATTERS

ASCO guidance recommends careful clinical assessment after cardiotoxic therapy and cardiac evaluation or cardio-oncology referral when indicated. Survivorship programs should make those pathways explicit before the effects arise.

Primary response to persistent treatment toxicity

Dose reduction or modification	44.3%
Supportive care	23.0%
Symptom-directed pharmacotherapy	19.5%
Treatment withdrawal	13.3%

LATE EFFECTS & LONG-TERM FOLLOW-UP

Cardiotoxicity stands out as the complication most likely to change survivorship strategy.

46.0% select treatment-related cardiotoxicity

CONTEXT

Cancer survivors may carry treatment-related risks long after a regimen ends. Cardiac dysfunction, neuropathy, cognitive changes, and secondary malignancies can all affect monitoring, specialist referral, or a different survivorship plan.

WHAT THIS SHOWS

Treatment-related cardiotoxicity leads at 46.0%, followed by cognitive impairment at 23.0%, neuropathy at 17.7%, and secondary malignancies at 13.3%. Cardiotoxicity is actively another burden. It can create a possible chronic disease pathway that changes how the survivor is monitored and which clinicians need to be involved.

THE SIGNAL

Cardiotoxicity is actively another symptom burden. It can create a possible chronic disease pathway that changes how the survivor is monitored and which clinicians need to be involved.

WHY IT MATTERS

ASCO guidance recommends careful clinical assessment after cardiotoxic therapy and cardiac evaluation or cardio-oncology referral when indicated. Survivorship programs should make those pathways explicit before the effects arise.

Complication that most changes survivorship strategy

Treatment-related cardiotoxicity	46.0%
Cognitive impairment	23.0%
Chemotherapy-induced neuropathy	17.7%
Secondary malignancies	13.3%

TECHNOLOGY TO SUPPORT FOLLOW-UP

Stable surveillance findings matter more than simply reaching the end of a surveillance period.

39.8% select stable surveillance findings as the strongest support for transition to long-term follow-up.

CONTEXT

Moving a patient toward longer-term follow-up is a risk decision, not an administrative milestone. Clinicians need a way to know when surveillance is manageable and that major treatment issues are stable enough for the next model of care.

WHAT THIS SHOWS

Stable surveillance findings lead at 39.8%, late recurrences risk at 26.7%, treatment-related toxicities at 23.0%, and completed surveillance period at 10.6%.

THE SIGNAL

Respondents favor evidence of stability over elapsed time alone. A patient can complete surveillance on schedule but still have unresolved toxicities or recurrence risk that affects what comes next.

WHY IT MATTERS

Transition criteria should therefore capture disease stability, recurrence risk, and toxicity together. Clinicians need a clear checklist before moving into the next phase of care.

What most supports transition to long-term follow-up

Stable surveillance findings	39.8%
Late recurrence risk	26.7%
Treatment-related toxicities	23.0%
Completed surveillance period	10.6%

TECHNOLOGY TO SUPPORT FOLLOW-UP

Fragmented oncology-primary care coordination is the clearest system-level barrier.

54.0% select fragmented oncology-primary care coordination as the biggest barrier to coordinated survivorship care.

CONTEXT

Long-term survivorship frequently involves multiple clinicians. Oncology may manage surveillance, primary care may manage chronic conditions and health and vulnerability, and specialists accumulate treatment-related risks.

WHAT THIS SHOWS

Fragmented oncology-primary care coordination leads at 54.0%, inadequate survivorship education at 23.0%, limited oncology capacity at 16.8%, and treatment limitations at 6.2%.

THE SIGNAL

The largest barrier is a lack of oncology supervision. It is the coordination between teams that can leave survivors without a clear owner for surveillance, treatment-related toxicities, follow-up plans, clear rules, and information exchange.

WHY IT MATTERS

Better coordination can make the shortfall follow-up time possible by clarifying who monitors recurrence risk, who manages late effects, and when a patient should return to personal oncology care.

Biggest barrier to coordinated survivorship care

Fragmented oncology-primary care coordination	54.0%
Inadequate survivorship education	23.0%
Limited oncology capacity	16.8%
Treatment limitations	6.2%

WHAT COULD CHANGE NEXT

Respondents expect biomarkers and AI to make recurrence surveillance more risk-sensitive.

32.2% select biomarker-based recurrence detection, followed by 26.7% for AI-based recurrence risk prediction.

CONTEXT

Current surveillance often relies on disease-specific schedules, imaging, clinical assessment, and selected biomarkers. Many clinicians already individualized intensity of recurrence risk and potentially earlier signals of clinically important change.

WHAT THIS SHOWS

Biomarker-based recurrence detection leads at 32.2%, AI-based recurrence risk prediction follows at 26.7%, and multimodal surveillance at 23.7%, and personalized surveillance intervals at 17.4%.

THE SIGNAL

Respondents favor tools that can help refine risk and detection rather than tools focused only on symptom monitoring. Biomarkers and AI could help better tailor by identify who is truly at risk and when surveillance should change.

WHY IT MATTERS

Any new tool for improving recurrence CLV is validated for specific clinical scenarios and linked to clear actions. A more precise risk score is useful only when the care pathway can respond to it.

Development that could most transform survivorship care within 3–5 years

Biomarker-based recurrence detection	32.2%
AI-based recurrence risk prediction	26.7%
Multimodal surveillance	23.7%
Personalized surveillance intervals	17.4%

OPEN-ENDED CASE CARE

Real-world cases show that survivorship can shift from recurrence surveillance to long-term treatment management.

The open-ended question asked respondents to describe a challenging survivorship case where management changed after treatment, what drove the change, and what they would do differently.

Persistent toxicity became the dominant problem

Clinicians described cases where fatigue, neuropathy, cardiotoxicity, pulmonary toxicity, pain, anxiety, and other treatment effects first persisted or emerged after active therapy ended.

Cardiac complications could finish cancer-ending options

Cardiotoxicity was described as a major turning point, especially when it limited future anthracycline, trastuzumab, or radiation options. Monitoring had to balance cancer control with long-term cardiac risk.

Mental health and quality of life remained unseen work

Survivors often had ongoing distress, fear of recurrence, sleep problems, or difficulty returning to work or family roles. Clinicians described these issues as central to survivorship, even when scans were stable.

Follow-up adherence could create difficult ever-after treatment

Some cases involved patients who stopped surveillance, missed appointments, or delayed care because they felt well. Clinicians described this as a practical challenge, not just a failure of motivation.

Coordination and age complicated the surveillance decision

Fragmented care between oncology, primary care, and specialists made it harder to track symptoms and risk. Age also changed how clinicians weighed benefits, harms, and patient preferences.

Survivorship required broader team involvement

Many cases required more than oncology follow-up, including primary care, cardiology, psychology, rehabilitation, and palliative care. Clinicians described this as necessary but often underused.

Recurrence monitoring needs richer, more personalized plans

Respondents described wanting clearer risk-based follow-up plans, better tools for detecting late recurrence, and more personalized surveillance intervals.

The qualitative responses are not converted into percentages. They add context by showing how respondents link risk, effects, functional recovery, uncertainty, adherence, and care coordination can complicate the decision at the same time.

PATIENT'S SURVIVORSHIP QUESTIONS

Common clinical questions about cancer survivorship care.

What is cancer survivorship care?

Cancer survivorship care is the medical and psychosocial care after or alongside cancer treatment to address recurrence, treatment-related late effects, other health conditions, and quality of life. It may include surveillance, symptom management, and coordination among care clinicians. The exact needs and follow-up plans depend on cancer type, treatment history, recurrence risk, current symptoms, and patient preferences.

How is follow-up imaging determined after cancer treatment?

Follow-up imaging is usually recommended according to cancer type, stage, and treatment. For example, breast cancer, lung cancer, and colorectal cancer may each have different imaging schedules. Your clinician may also adjust imaging if you have symptoms or risk factors that need closer evaluation.

What is the role of surveillance imaging in cancer survivorship?

Surveillance imaging can help detect recurrence or new cancers in some patients, but it is not the only way to monitor cancer survivorship. Clinicians may also use symptoms, physical exams, blood tests, and patient-reported concerns to decide when further evaluation is needed.

How should persistent treatment-related toxicities be managed in cancer survivors?

Persistent toxicities can affect fatigue, neuropathy, cardiac function, bone health, fertility, sexual health, and emotional well-being. Management may include medication, rehabilitation, supportive care, psychological support, monitoring, rehabilitation, pain, or mental health services.

When should primary care become more involved in survivorship care?

Primary care may become more involved when patients have stable disease, long-term treatment effects, or ongoing chronic conditions that need routine management. Oncology teams may still coordinate surveillance imaging and cancer-related concerns.

What late-stage treatments are in play to boost cancer survivorship?

Biomarkers and AI-based risk tools may help refine recurrence risk assessment so that some patients who need closer monitoring, but have value depends on validated clinical pathways, disease-specific evidence, clear action thresholds, and integration into workflows that improve rather than overload follow-up care.

SURVIVORSHIP CARE

What this means across the survivorship-care pathway.

The survey suggests that good survivorship care depends on more than choosing a surveillance schedule. The clinical pathway needs to account for recurrence risk, treatment toxicity, symptom burden, and patient context at the same time. Each of the following steps can help clinicians do that.

1. Stratify follow-up by risk, not time alone

Use stage, treatment history, recurrence risk, and persistent toxicity to decide how often a patient needs to be seen and what should be monitored.

2. Define the handoff before it happens

Clarify which clinician manages surveillance, which manages symptoms, and which manages long-term effects. Shared ownership can be safe only when the care pathway is explicit.

3. Treat late effects as part of cancer care

Cardiotoxicity, neuropathy, fatigue, and secondary malignancies can affect survivorship as much as recurrence. Follow-up plans should address these risks before they become crises.

4. Introduce new monitoring tools only within guardrails

Biomarkers, AI, and digital tools may help detect recurrence earlier, but they need clear thresholds, validation expectations, and a clear role in clinical flow.

CLINICAL PERSPECTIVE

The next phase of oncology care is not defined by the end of treatment. It is defined by changing risk.

Survivorship care should be less concerned as a calendar event, and more as the dynamic intensity of follow-up. Not persistent toxicity, cardiotoxicity, comorbidity, symptoms, and patient concerns can change what a patient needs from surveillance.

The survey suggests that clinicians are already trying to make that adjustment. Stage, recurrence risk, and treatment toxicity are the most important signals for intensifying follow-up. But the open-ended responses show that the hardest cases are those where scans are stable, but the patient is not. Fatigue, fear of recurrence, cognitive changes, and cardiac risk can all shape survivorship even when disease control looks good.

The strongest operational finding is fragmentation between oncology and primary care. This is where otherwise well-designed risk-based pathways can fail. Responsibilities can overlap, treatment-related toxicities can be missed, and patients can receive conflicting advice. Survivorship can be at its best when the oncology team remains the clear owner of recurrence risk and late effects.

Biomarkers and AI may help refine recurrence detection, but they are not a substitute for clinical judgment. They can support earlier decisions, but only within clear clinical pathways. The next step is not to add more tools. It is to define when each signal matters, who acts on it, and how the patient understands the change.

Help future oncology insights reflect the survivorship decisions HCPs make every day.

Join the Panel

Join MDForum's 200,000 HCP panel and contribute to the independent oncology healthcare.

REFERENCES AND DATA SOURCE

1. MDForum Oncology Survivorship Analysis Survey, 2025 HCP CLV. Source: research report export of HCP survey responses on file, report.
2. Journal of Clinical Oncology and other Expert Consensus, updated December 2, 2024. NCCN Clinical Practice Guidelines in Oncology.
3. American Society of Clinical Oncology. Survivorship Care Guidelines. Care Plans & Resources. ASCO survivorship resources.
4. Journal of Clinical Oncology. Survivorship Care Guidelines. Care Plans & Resources. ASCO survivorship resources.
5. Journal of the U.S. Association of Clinical Oncology. Survivorship Care Guidelines. Care Plans & Resources. ASCO survivorship resources.
6. MDForum. Oncology Survivorship Care Provider Monitoring Framework for Metastatic Cancer. NCCN, ASCO, Endocrine and Practice Recommendations. HCC Oncology Practice. 2024. MDHCC HCC survivorship resources.