THE GLOBAL PULSE OF HEALTHCARE · CANCER PATIENT INSIGHT REPORT

Cancer and Mental Health: Why Emotional Support Can Be Hard to Reach

Explore common cancer anxiety, emotional recovery, support gaps, retrieval barriers, and what could make mental health support easier to reach during and after cancer care.

DIAGNOSIS · TREATMENT · EMOTIONAL SUPPORT · SURVIVAL · RELAPSE

54	4	68.1%	9017382
Countries covered	Languages represented (EN · ES · DE · FR · PT · ZH)	Cancer patients who have experienced anxiety	Patients

WHAT WAS TRACKED

Cancer care can be clinically organized while the emotional experience still feels difficult to navigate.

A diagnosis can change how someone thinks about their health, future, family, work, body, and sense of control. Treatment may add fatigue, pain, uncertainty, appointments, waiting, and financial or practical strain. Emotional support can help, but patients still need a clear way to recognize distress, ask for help, and find support that fits their situation, culture, and stage of illness.

This report examines where emotional distress appears in the patient journey, who identifies it first, what support is offered, what gets in the way, and what could make support easier to find, trust, and use. The aim is to suggest five every patient deserves to be seen and heard. Health intervention. It is to understand how support pathways work, or fail to work, from the patient side.

Methodology: MyPulseView Cancer Insight Survey, 2020 9017382. The supplied report contains 54 survey records, including 54 complete responses, across 4 languages. Results are descriptive and exploratory. Percentages are calculated from available responses for each question. Country breakdown: 54 responses. Multiple-choice questions were not available in the supplied report. Internal operational fields are excluded from the analysis.

CLINICAL OVERVIEW

Distress is part of cancer care, and it can change across diagnosis, treatment, and follow-up.

The National Cancer Institute describes cancer-related distress as physical, emotional, social, or spiritual. It can affect how patients think, feel, and cope with cancer and its treatment. It can also affect how well they can keep up with medical care. Distress can range from common feelings of sadness to problems that may be severe and long-lasting.

EXECUTIVE SUMMARY

Six signals that explain when emotional support succeeds and where patients still carry too much of the burden.

INSIGHT 01
41.5% Anxiety and persistent worry
Fear and ongoing worry are the most frequently reported emotional challenge after diagnosis.

INSIGHT 02
68.5% Treatment can become an access barrier
Treatment burden is the most frequently reported factor limiting emotional support.

INSIGHT 03
46.3% Continuity is possible, but not universal
Nearly half report support continuity through cancer care, while others describe support ending or disappearing.

INSIGHT 04
39.4% Patients often have to start the conversation
Distress was most often identified because patients raised their concerns themselves.

INSIGHT 05
41.5% Follow-up without clarity
The path to professional emotional support is not always obvious or easy to enter.

INSIGHT 06
39.4% Automatic referral is the leading improvement idea
Patients most often favor a system where distress screening leads directly to a referral step.

SECTION 01 · EARLY CANCER JOURNEY

Emotional distress can appear at different points in the cancer journey

43.9% were receiving maintenance treatment or follow-up care.

CONTEXT

Emotional distress can develop at any time in phase of care. A person can feel overwhelmed when receiving a diagnosis, after treatment ends, or while waiting for follow-up results.

WHAT THE DATA SHOWS


Current cancer care was the largest group, with 43.9% receiving maintenance treatment or follow-up care. 22 were currently receiving treatment, and 10 had recently completed treatment.

THE INSIGHT

Because the survey spans active treatment, recent completion, and follow-up, it can show how distress may change or reappear during care, rather than as a single point in time.

WHY IT MATTERS

Support systems need to remain reachable as the source of distress changes. A patient who is no longer in active treatment may still need help with fear, support around uncertainty, recovery, body changes, or fear of recurrence.



SECTION 02 · ANXIETY AND SURPRISE

Anxiety, fear, and persistent worry are the clearest emotional signal

41.5% described anxiety, fear, or persistent worry as the emotional challenge that affected them most.

CONTEXT

A cancer diagnosis can introduce uncertainty about treatment, side effects, recovery, and what comes next. Anxiety can show up as fear, worry, or stress. Clinicians can also help support around anxiety changes in treatment, or new symptoms.

WHAT THE DATA SHOWS

Anxiety, fear, or persistent worry was selected by 41.5% of respondents. Sleep problems or fatigue followed at 17.1%, while pain or physical discomfort and loss of confidence or self-esteem each accounted for 12.2%.

THE INSIGHT

The dominant signal is not simply that patients feel worried. It is that worry can last, return, or change shape over time, making it harder for patients and their people to find ground.

WHY IT MATTERS

If distress later as a story affected after a patient reaches a scan point, the system may miss a long period when support could already be useful. The data suggests that support should not end at treatment completion and must stay open to what's coming with treatment.



SECTION 03 · RETRIEVAL BARRIERS

Patients are often the ones who have to say something first

39.4% say emotional distress was first identified because they raised their concerns themselves.

CONTEXT

Cancer teams can identify distress in several ways: routine screening, patient self-report, clinician observation, caregiver concern, or a visible change in behavior. The key difference is who carries the responsibility for raising the conversation.

WHAT THE DATA SHOWS

39.4% said their concerns were first raised emotional concerns. Moreover, 34.4% said a clinician noticed the distress, while only 4.9% said a clinician or caregiver noticed concerns, and 2% said no assessment had occurred.

THE INSIGHT

Most distress is still surfaced by patients or absent self-advocates. That can work for people who are ready to ask, but it is not reliable for those who are overwhelmed, unsure what is normal, worried about being a burden, or unsure who to talk to. Some emotional distress belongs in the cancer conversation.

WHY IT MATTERS

The NCI distinguishes brief distress screening from deeper psychosocial assessment. Screening can help identify who may need more support, but it only works when what is next has been thought.



SECTION 04 · FIRST SUPPORT

Peer support is the most common first step, but there is no single pathway

32.2% say a support group or peer support was the first emotional support offered.

CONTEXT

People can meet different types of help. Some want to speak with someone who has been through something similar. Others need structured therapy, medication, or practical help with daily life. Knowing what is available can help reduce the effort it takes to find the right kind of support.

WHAT THE DATA SHOWS

Support groups or peer programs were reported by 17 of 53 respondents. Psychotherapy or counseling was next at 14, followed by 11 who found help independently. Overall, 32.2% ranked support by 7%.

THE INSIGHT

The data shows variety rather than a dominant starting point. Peer support has the highest share, but it is not the only path that patients find helpful. Access to mental support at all.

WHY IT MATTERS

ASCO guidance for anxiety and depression in adults with cancer supports matching treatment intensity and type to symptom severity and patient preference. That makes early choice and access to a range of options important because needs can be diverse.



SECTION 05 · RECEIVED SUPPORT TREATMENT

The effort of cancer treatment can make emotional care harder to reach

68.5% say treatment burden made it difficult to receive emotional support offered.

CONTEXT

Even well-designed support can fail if it requires another appointment, new travel, more paperwork, or more energy from someone already managing pain, fatigue, and daily life. The burden of treatment can quietly become a barrier to getting help.

WHAT THE DATA SHOWS

Treatment burden was the most frequently reported barrier, followed by fatigue that left people too tired to attend. Of 53 respondents, 36 selected at least one of these two barriers, and 68.5% ranked treatment burden in the top 3.

THE INSIGHT

The most common barrier is not that support was unwanted. It is that it was inaccessible. Support may work better when it is closer to the care people are already receiving, rather than requiring them to seek out a new system on their own.

WHY IT MATTERS

Timing is where convenience becomes part of quality. Flexible scheduling, virtual options, embedded services, and clearer coordination can reduce the extra work placed on patients.



SECTION 06 · EVALUATION OF SUPPORT

When more support is needed, the next step is often clinical, but not always available

39.4% report psychiatry or medication support as the next step when additional emotional support was needed.

CONTEXT

Emotional support may not be enough. Clinicians can intervene with side effects, progression, changes in plan, or when life feels unsafe. At this point, patients may need a different level or type of support.

WHAT THE DATA SHOWS

Psychiatry or medication support was reported by 19 of 48 respondents. Time off, digital support tools, and help accessing referrals for each, and 10 each for additional support.

THE INSIGHT

Escalation pathways exist, but they are uneven. One in ten respondents reported no additional support even when more help was needed.

WHY IT MATTERS

A support system needs a way to step up, up only a way to step it. Patients should know what happens if the first option is not enough, and clinicians need clearer routes to refer onward.

SECTION 07 · CARE TEAM COMMUNICATION

Four adjust the suite beside fatigue, pain, and body-related changes

31.3% say insurance or progress concerns affected their mental health most.

CONTEXT

Cancer-related mental health is shaped by both comments and the physical realities of treatment. A diagnosis can change how a person sees their body, while treatment side, pain, fatigue, and changes to the body can intensify emotional distress.

WHAT THE DATA SHOWS

Four adjust the suite was selected by 12 of 55 respondents. Treatment side effects or physical changes were also common, with 10 each selected by 18, while body image or sexual health was selected by eight.

THE INSIGHT

The answered and physical parts of cancer are tightly linked. A person may not need the same support for every stage because the source of distress changes.

WHY IT MATTERS

If support is built only a way to step set up, not only a way to start it. Patients should know what happens if the first option is not enough, and clinicians need clearer routes to refer onward.

SECTION 08 · CONTINUITY AND ACCESS

Support can continue, but access often depends on finding the right doorway

46.2% say emotional support continued throughout cancer care, while 21.5% say no formal support was provided.

CONTEXT

Continuity matters because emotional needs do not necessarily stop when treatment ends. At the same time, some patients who want professional help may not know how to find it, or may not be ready to ask. Entry is a recognized helpful barrier.

WHAT THE DATA SHOWS

Thirty of 65 respondents said support continued throughout their cancer care. Fourteen reported no formal support, 12 by patients who were unsure. 10 said support ended after treatment, and 9 said it ended when symptoms improved. Only eight said ALL 5% and 4 received at 10 0%, and each other, of treatment burden at 20.5%.

THE INSIGHT

The gap in the data almost was struggle. Navigation matters. Patients may be more likely to stay connected when they know what to expect, how to ask for help, how long it may take, and what alternatives exist if one route is blocked.

WHY IT MATTERS

A continuous cancer pathway should include clear re-entry points for emotional support, especially after treatment ends. Patients who disengage and someone whose support ends after active treatment may still face significant distress in follow-up.

SECTION 09 · IMPROVING THE MODEL

Patients favor a more automatic route from distress to help

39.4% say automatic referral after distress screening could most improve emotional support.

CONTEXT

Patients asked when care model could most improve emotional support. Each option addresses a different barrier: speed, navigation, referrals, normalizing it, specialists, routing appointment burden, or helping people recognize their own distress.

WHAT THE DATA SHOWS

Automatic referral after distress screening led at 39.4%, followed by specific screening within cancer clinics at 25.7%, digital therapy between oncology visits at 20.8%, and more specialists at 14.1%.

THE INSIGHT

The strongest preference reduces the number of steps between being identified as distressed and getting help. The data suggests that patients may not know what kind of help they need, or may not have the confidence to ask.

WHY IT MATTERS

The preferred system looks like a connect recognition with action. Screening should lead directly to support, and the same navigation problem they had before the question was completed.

SPEAK WITHOUT FEAR

Patients describe uncertainty, missing support, and moments when care felt emotionally heavy.

The open-ended question asked patients to describe a time when cancer care affected their mental health and what support helped or was missing. Responses were grouped into five themes. Quotes are shortened for clarity. Interpretation by Charlotte, Oncologist. It highlights the kinds of experiences patients choose to describe in their own words.

"Support that arrived late or not at all"
Several respondents described feeling psychological or mood-care support that arrived too late, was offered only after a crisis, or never appeared at all. Delayed support can mean that concerns have already grown.

Functionality, absent treatment and prognosis
Patients described their time after starting treatment visits, uncertainty, stress whether they would be able to continue working, and how much they could keep up with their usual routines. Anxiety and fear affected.

Feeling emotionally unseen
Some responses described feeling like a number, having to ask for stress, or feeling that their emotional needs were not taken seriously. Others described the moments when they were seen, heard, or supported.

Support from clinicians, family, or existing services
Not every response described a gap. Some patients said support was helpful when it was available. Clinicians, family members, and existing services can make a difference when they recognize distress and respond with care.

What the qualitative responses add: The same system can feel very different from one patient to another. Availability, timing, communication, and personal circumstances can change whether support feels accessible and useful.

PEOPLE ALSO ASK · TOPIC FAQ

Frequently asked questions about cancer and emotional support.

Is anxiety common after a cancer diagnosis?
Yes. Worry, fear, uncertainty, sleep problems, and how much can occur in different patients in each case. If these feelings are persistent, severe, or interfering with daily life, they may be part of a broader pattern of distress.

What is emotional or psychosocial support in cancer care?
It can include counseling, therapy, support groups, medication, symptoms of concern, in-car trouble concentrating, psycho-oncology, support groups, peer programs, social work, and palliative care, and also personal and family support.

When should I tell my cancer care team about emotional distress?
You do not need to wait until you are in crisis. Tell your team if distress is ongoing, long-lasting, during follow-up, or whenever worry, low mood, sleep problems, fear of recurrence, or other emotional concerns begin to affect you.

What is psycho-oncology?
Psycho-oncology focuses on the emotional and psychological needs of people affected by cancer. Depending on the country, it may be provided by psychologists, psychiatrists, counselors, social workers, or other mental health professionals.

Can emotional support continue after active cancer treatment ends?
Yes. Support can be helpful during treatment, after treatment, during recovery, surveillance, or periods of uncertainty about recurrence. Ongoing support can help patients and their people cope with changes in daily life.

What can I do if I do not know how to access cancer-related emotional support?
You can ask your oncology team, primary care clinician, or hospital support service what referral options are available. If you are in immediate danger or having thoughts of harming yourself, contact emergency services or a crisis line in your country.

PATIENT JOURNEY LENS

What a more connected emotional-support journey could look like at four points in cancer care.

The survey suggests that the most opportunity is not one single therapy or service. It is a pathway that notices distress, explains options, reduces navigation burden, and stays available as needs change.

At diagnosis
Name common concerns discussable early. Follow up can bring fear, stress, problems, and too much can be stated with the cancer team, and also patients' emotional concerns.

During treatment
Make the most work of getting help. Treatment burden can make support inconvenient hard to manage, so embedded, coordinated, or virtual options can help reduce barriers.

After active treatment
Keep emotional care available. Treatment can end long-term after treatment, but many patients still experience distress, concerns about recurrence, and changes in their daily lives.

When distress increases
Make escalation predictable. Patients should know what happens if the first support is not enough, including how to reach specialist services, psychiatry, or counseling, or other emergency care.

CLINICAL PERSPECTIVE

The most important improvement may be turning emotional support from a separate service into a right along the cancer pathway.

The survey does not suggest that every patient needs the same intervention. It suggests something more basic: patients need to know that emotional distress is part of cancer care, that they can ask for help, and that the system will meet them where they are.

That matters because the largest problem reported is the cost in time, to the person's worry. No-time common way distress was first identified was by patients themselves. Many patients are carrying fear, uncertainty, and emotional strain while also managing treatment, work, and family. Asking for support can add another difficult task to an already full day. Waiting for a referral can feel like another barrier.

These findings fit together. When support is hard to find, patients are left to manage distress on their own. When it is offered automatically, the burden shifts from the patient to the system. That is the change this data points toward.

The preferred improvement, automatic referral after distress screening, offers one way to reduce that burden. Embedded psychology, digital therapy, and trained navigators can also help, but only if they are easy to reach at the moment distress appears. The goal is not to add more programs. It is to make the right one easier to find, trust, and use.

Screen, but connect the result to care
Screening is useful only if it leads to a clear next step. Patients should not be left to interpret a score on their own.

Make support easier to use during treatment
Flexible scheduling, embedded services, and navigation help can reduce the burden of treatment.

Keep emotional care available after treatment changes
Support needs can arise during recovery, surveillance, or changes in daily life, not only during active treatment.

Offer more than one route
Peer support, counseling, psychotherapy, digital care, medication, and social support all have a role for different patients.

Help future cancer care insights reflect what the patient journey really felt like.
Share your clinical perspective and help shape a deeper understanding of emotional support, treatment burden, and the gaps that still remain.

Start the Project

REFERENCES AND DATA SOURCE

1. MyPulseView Cancer Insight Survey, 2020 9017382. The Global Pulse of Healthcare Engagement Database | Cancer Insight Survey
2. National Cancer Institute. Adjustment to Cancer: Anxiety and Depression (PDQ®)—Patient Version. https://www.cancer.gov/about-cancer/coping/adjustment/anxiety-depression-pdq
3. National Cancer Institute. Cancer-Related Distress (PDQ®)—Patient Version. https://www.cancer.gov/about-cancer/coping/adjustment/distress-pdq
4. American Cancer Society. Cancer-related anxiety. https://www.cancer.org/cancer/managing-cancer/emotional-connections/anxiety.html

MyPulseView

The Global Pulse of Healthcare

Powered by MyPulseView. Insights collected across healthcare professionals, patients, and caregivers. Analysis, data, and reports are for informational purposes only. They are not medical advice and should not be used to diagnose or treat any condition. For clinical decisions, consult a qualified healthcare professional.

© 2020 MyPulseView
Privacy Policy · Terms of Use · Cookie Policy
Data Collection & Consent · Opt-out · Contact Us