Complex Diabetes Care: When HbA1c Is Only Part of the Decision

Insights and professional perspectives on glycemic control, obesity, kidney and cardiovascular risk, treatment sequencing, adherence, access, and sustained therapy value.

EDITORIAL STRATEGY · COMMENTARY PIECE · THERAPY CHOICE · LONG-TERM OUTCOMES

46	8	65.6%	9001530
Pages	Expert contributors from UK, Spain, Italy, France, Germany	Adults with diabetes	ISBN

Survey theme: 46 survey records with a 65.6% completion rate across the USA, UK, Canada, Italy, France, and Germany. Substantive question biases range from 31 to 46 responses. Internal reproducibility: maximum bias, split-sample, and Kolmogorov tests consistent from analysis.

CLINICAL OVERVIEW

Current diabetes guidance increasingly treats therapy choice as a risk-management decision, not only a glucose-lowering decision.

The ADA 2026 Standards of Care position that glycemic control remains the foundation of diabetes management, but with cardiovascular and kidney disease, weight, hypoglycemia risk, cost, and access as decisive factors, effectiveness, tolerability, and individual preferences.

For adults with type 2 diabetes and established or high risk for cardiovascular risk, these same guidance places SGLT2 inhibitors and/or GLP-1 receptor agonists with demonstrated benefit directly into the treatment plan, in appropriate patients, irrespective of the metformin or baseline glycemic levels. The ADA section also recommends routine assessment of urine albumin and estimated glucose, and the obesity section positions weight management as a central component of diabetes care, not a separate concern.

Taken together, these updates reflect a shift toward more individualized, risk-based therapy. The follow-up MASTLocate survey findings will show to what degree this is reflected in real-world clinical practice, and to what extent it is already being implemented.

EXECUTIVE SUMMARY

Six findings that show why complex diabetes care increasingly extends beyond HbA1c.

01

57.9%

Obesity leads the comorbidities challenge

Obesity is the problem most often selected as complicating diabetes management.

02

55.3%

HbA1c still anchors decisions

Most clinicians still use HbA1c as the primary tool to choose the most advanced assessment influencing treatment choice.

03

44.4%

SGLT2 inhibition leads the next step

Among advanced options, adding an SGLT2 inhibitor is the most selected next action.

04

42.9%

Results can extend beyond HbA1c

Cardiovascular risk reduction is the most valued reason to continue therapy despite suboptimal HbA1c.

05

45.7%

Adherence is the biggest complex-care barrier

Poor adherence, limited affordability, access, and fragmented care as the selected barrier.

06

35.3%

Earlier cardiorenal direction leads future

Earlier cardiorenal risk direction is the most selected future development by the development that could transform care.

01 · PERSPECTIVE PIECE

Obesity is the comorbidity most often selected as complicating diabetes management

57.9% select obesity, compared with 31.4% for chronic kidney disease and 19.5% for cardiovascular disease.

CONTEXT

Obesity can change glycemic targets, weight goals, insulin requirements, hypoglycemia risk, and the way the patient and the provider discuss treatment intensification and long-term safety.

WHAT THE DATA SHOWS

Obesity leads the audience at 57.9%, CKD follows at 31.4%, while cardiovascular disease accounts for 19.5%.

WHY IT MATTERS

The ADA 2026 obesity guidance positions weight management as a primary therapeutic objective, not a secondary concern. Specialists are aligned with this: type 2 diabetes and overweight is already the default.

THE HEADLINE

Obesity is already the defining signal, but most clinics is already tentatively influencing treatment decisions for a substantial share of respondents.

WHY IT MATTERS

Current ADA 2026 guidance makes kidney assessment clinically consequential. Underdiagnosis between extremes and large protective gains over therapy with SGLT2 and eGFR.

Comorbidity that most complicates diabetes management

Obesity	57.9%
Chronic kidney disease	31.4%
Cardiovascular disease	19.5%

02 · DECISION DRIVER

HbA1c still anchors treatment decisions, but renal markers are already part of the decision frame

55.3% select HbA1c and the glucose profile, 23.7% select eGFR and albuminuria.

CONTEXT

Kidney function is a familiar measure of glycemic exposure, yet complex diabetes care increasingly requires it to be read alongside albuminuria, weight, cardiovascular risk, and treatment safety alongside the glucose profile.

WHAT THE DATA SHOWS

HbA1c and the glucose profile lead at 55.3%, followed by eGFR and albuminuria at 23.7%, weight and metabolic profile at 15.8%, and ASCVD risk at 5.3%.

THE HEADLINE

Most clinicians describe starting signal, but renal status is already tentatively influencing treatment decisions for a substantial share of respondents.

WHY IT MATTERS

Current ADA 2026 guidance makes kidney assessment clinically consequential. Underdiagnosis between extremes and large protective gains over therapy with SGLT2 and eGFR.

Assessment that most influences treatment decisions

HbA1c and glucose profile	55.3%
eGFR and albuminuria	23.7%
Weight and metabolic profile	15.8%
ASCVD risk assessment	5.3%

03 · CLINICAL THERAPEUTICS

The first treatment priority is split rather than dominated by one default approach

Metformin-based therapy and GLP-1 receptor agonists are tied at 23.3% each.

CONTEXT

Modern treatment selection depends on more than glucose-lowering potency. SGLT2 inhibitors, cardiovascular disease, hypoglycemia, tolerability, access, and patient preferences can all change what makes sense first.

WHAT THE DATA SHOWS

Metformin-based therapy and GLP-1 receptor agonist each receive 23.3%. SGLT2 inhibitors and dual GIP/GLP-1 therapy each receive 16.7%.

THE HEADLINE

The result right suggests the respondents are not operating from one universal first-line sequence. The starting point appears sensitive to the clinical profile that is being considered.

WHY IT MATTERS

Current ADA 2026 therapeutic guidance similarly emphasizes person-centered choice, including comorbidities, weight, hypoglycemia, cost, adverse effects, and preferences.

Therapy prioritized initially

Metformin	23.3%
GLP-1 RA	23.3%
SGLT2i	16.7%
GIP/GLP-1	16.7%

04 · ACCESS & ECONOMICS

Cost and access limit treatment optimization more often than treatment effectiveness

39.4% select access as the leading limitation, followed by adverse effects or tolerability at 30.8%.

CONTEXT

Clinical choice only becomes meaningful when the patient can obtain the medication, tolerate it, understand how to use it, and continue it over time.

WHAT THE DATA SHOWS

Access is ranked leader at 39.4%, tolerance effects or tolerability follow at 30.8%, hypoglycemia risk at 15.4%, and complexity or burden at 14.4%.

THE HEADLINE

The survey places practical access and tolerability concerns ahead of glycemic complexity as optimization barriers. The gap between preference and favorable care may therefore emerge before the biology is compromised.

WHY IT MATTERS

ADA 2026 guidance explicitly recommends assessing financial obstacles and working across the care team to reduce cost-related barriers where possible.

What most limits treatment optimization

Cost or access	39.4%
Adverse effects or tolerability	30.8%
Hypoglycemia risk	15.4%
Complexity or burden	14.4%
Treatment uncertainty	11.1%

05 · DRUG CLASS DISCUSSION

After inadequate glycemic control, SGLT2 inhibition is the most selected next step

44.4% prefer adding an SGLT2 inhibitor; 22.2% prefer adding a GLP-1 receptor agonist.

CONTEXT

When control remains inadequate, HCPs can intensify the current regimen, move toward insulin, or add a class they have not used before. Preference may not always match guideline order of the treatment algorithm.

WHAT THE DATA SHOWS

Adding an SGLT2 inhibitor tops the list at 44.4%, followed by GLP-1 receptor agonists at 22.2%. Intensifying existing therapy and switching insulin each account for 11.1%.

THE HEADLINE

The preferred next step shifts toward newer classes with potential benefits beyond glycemia rather than simply escalating the current backbone.

WHY IT MATTERS

The ADA's pentagonal care model emphasizes person-centered choice, including comorbidities, weight, hypoglycemia, cost, adverse effects, and preferences.

Preferred next step after inadequate glycemic control

Add SGLT2i	44.4%
Add GLP-1 RA	22.2%
Intensify current therapy	11.1%
Switch to insulin	11.1%
Other	11.1%

06 · DRUG CLASS DISCUSSION

Advanced CKD and heart failure are the clearest reasons to change treatment strategy

34.3% select advanced CKD, followed by heart failure at 25.7%.

CONTEXT

Comorbidity can change medication choice, dosing, safety, expected benefit, and monitoring. Kidney disease, heart failure, and heart fibrosis, shape protection decisions point of the treatment objective.

WHAT THE DATA SHOWS

Advanced CKD leads at 34.3%, heart failure at 25.7%, while hypoglycemia risk at 20.0% and heart fibrosis at 20.0%.

THE HEADLINE

The results from kidney status appear to exert more influence on treatment sequencing than other listed subgroups.

WHY IT MATTERS

The ADA's cardiovascular and CKD sections connect therapy selection directly to comorbidities and kidney monitoring in stage-based guidance.

Subgroup that most changes treatment strategy

Advanced CKD	34.3%
Heart failure	25.7%
Established ASCVD	20.0%
Frail older adults	20.0%

07 · PATIENT PERSPECTIVE & ADHERENCE

Cardiovascular risk reduction is the leading reason to continue therapy despite suboptimal HbA1c

42.9% choose cardiovascular risk reduction, 28.6% choose weight reduction and 28.6% kidney protection.

CONTEXT

A treatment may be doing something clinically important even when glycemia alone does not look fully optimized. This question tests whether HCPs describe whether to maintain that therapy in order to benefit while aiming for improving gap elsewhere.

WHAT THE DATA SHOWS

Cardiovascular risk reduction leads at 42.9%, weight reduction at 28.6%, kidney protection at 28.6%, and hypoglycemia avoidance at 14.3%.

THE HEADLINE

Respondents are explicitly valuing outcomes beyond HbA1c, with cardiovascular benefit most likely to justify continuing a therapy.

WHY IT MATTERS

This aligns with current guideline logic in which cardiovascular and kidney benefits can influence therapy choice irrespective of the need for additional glucose lowering.

Outcomes that justify continuing therapy despite suboptimal HbA1c

Cardiovascular risk reduction	42.9%
Weight reduction	28.6%
Kidney protection	28.6%
Hypoglycemia avoidance	14.3%

08 · CARE DELIVERY

Poor treatment adherence is the biggest selected barrier to complex diabetes care

45.7% select poor treatment adherence, medication affordability follows at 34.3%.

CONTEXT

Complex diabetes care often asks patients to manage multiple medications, monitoring, lifestyle changes, appointments, side effects, and competing health priorities. Every additional requirement can create another point where the plan does not continue.

WHAT THE DATA SHOWS

Poor treatment adherence leads at 45.7%, followed affordability follows at 34.3%, fragmented multidisciplinary care at 11.4%, and limited specialist access at 8.6%.

THE HEADLINE

Adherence is the clearest lens a shortage of treatment options. It is surprising that behavioral gaps is real life.

WHY IT MATTERS

The ADA 2026 Systems of care guidance emphasizes person-centered team-based care, coordination, and integrated long-term management of diseases and comorbidities.

Biggest barrier to complex diabetes care

Poor treatment adherence	45.7%
Medication affordability	34.3%
Fragmented multidisciplinary care	11.4%
Limited specialist access	8.6%

09 · WHAT CHANGED NEXT

Earlier cardiorenal risk direction is the most selected development for transforming complex diabetes care

35.3% select earlier cardiorenal risk direction, followed by more effective combination therapies at 29.4%.

CONTEXT

The earlier kidney and cardiovascular risk becomes visible, the earlier treatment can be organized around protection rather than waiting for the complication to dominate the clinical picture.

WHAT THE DATA SHOWS

Earlier cardiorenal risk direction leads at 35.3%. More effective combination therapies follow at 29.4%, while more personalized treatment algorithms at 23.5% and personalized treatment algorithms at 11.8%.

THE HEADLINE

Respondents prioritize earlier recognition of risk over in purely algorithmic terms. The most selected future development is not a new drug but a shift toward of personalized algorithms.

WHY IT MATTERS

This future need suggested by the survey is not simply a more powerful drug. It is a system that identifies risk earlier, selects the right combination, and makes multidisciplinary care easier to execute.

Development that could most transform complex diabetes care

Earlier cardiorenal risk direction	35.3%
More effective combination therapies	29.4%
More personalized treatment algorithms	23.5%
Personalized treatment algorithms	11.8%

OPEN RESPONSE LENS

When HCPs describe complex cases, competing risks change the treatment plan before they change the goal.

The open-ended question asked respondents to describe a complex diabetes case in which competing risks changed their treatment strategy, what drove the decision, and what management would look like.

Renal function frequently changed the sequence

Respondents described CKD, albuminuria, high creatinine, advanced renal disease, and declining eGFR as the point at which the plan had to change.

Cardiorenal benefits were prioritized over glycemic only view

Several respondents described selecting or adding SGLT2 inhibitors or GLP-1 receptor agonists because cardiovascular, kidney, or heart failure risk was the central objective.

Weight and affordability created active trade-offs

Obesity, insulin-related weight gain, gastrointestinal intolerance, and the need to balance glucose control against hypoglycemia and quality of life were recurring themes.

Hypoglycemia and treatment burden complicated simplification

Respondents mentioned hypoglycemia risk, insulin management, polypharmacy, and patient burden as reasons they could not simplify the regimen as much as they would have preferred.

Affordability and adherence could override the plan

Medication cost and adherence appeared as practical constraints, sometimes preventing the use of the preferred agent even when it was clinically appropriate.

To single trade-off dominated every case

Some respondents prioritized glycemia, others kidney or cardiovascular risk, and others weight or tolerability. The most complex cases were those in which no single goal could be optimized without compromising another.

These themes are qualitative summaries of usable open-ended responses. They were not formally coded like percentages.

PEOPLE ALSO ASK · FAQ TABLE

Frequently asked questions about diabetes care and treatment decisions

How should chronic kidney disease change diabetes treatment choices?

Kidney function may change glycemic targets, medication choice, monitoring, and dosing. ADA 2026 guidance recommends therapies with demonstrated kidney and cardiovascular benefit for people with type 2 diabetes and CKD, while also accounting for eGFR, hyperkalemia risk, tolerability, and treatment burden.

When are SGLT2 inhibitors preferred in type 2 diabetes?

SGLT2 inhibitors are preferred when cardiovascular disease, heart failure, or cardiovascular risk makes cardiorenal protection a priority. Their use should reflect individual patient factors, including eGFR, albuminuria, volume status, and tolerability.

When are GLP-1 receptor agonists recommended in complex diabetes care?

GLP-1 receptor agonists are recommended when cardiovascular disease, cardiovascular risk reduction, or kidney-related benefit is relevant. Selection should still account for tolerability, contraindications, cost, and patient preferences.

Why might a diabetes therapy be continued when HbA1c is still above target?

If treatment can provide clinically meaningful benefits beyond glucose lowering. Depending on the therapy and the individual, cardiovascular risk reduction, kidney protection, weight reduction, or reduced hypoglycemia may justify continuing a therapy even when HbA1c remains suboptimal.

How does obesity influence diabetes treatment decisions?

Obesity can affect glycemic control, insulin resistance, and the choice of treatment. Current guidance recommends considering therapies that support weight reduction or avoid weight gain, especially when obesity is a major clinical concern.

What should healthcare professionals consider when cost or adherence limits treatment optimization?

Treatment plans need to be clinically effective and realistically sustained. When cost, access, tolerability, or adherence is limiting, care teams can prioritize therapies that are easier to obtain, better tolerated, more flexible, or more likely to be continued over time.

CLINICIAN DISAGREEMENT CLOUD

What this means at four points in a treatment decision.

The number suggests that HCPs describe complex diabetes cases in which competing risks changed their treatment strategy, what drove the decision, and what management would look like.

1. Open with the decision risk profile

Review glycemic burden, kidney function, albuminuria, cardiovascular disease, heart failure, weight, hypoglycemia risk, and treatment burden before choosing a therapy.

2. Define what the therapy is meant to protect

Be explicit about whether the main objective is glucose lowering, cardiovascular protection, kidney protection, weight reduction, or reduced hypoglycemia risk.

3. Surface kidney-driven monitoring implications

Identify when eGFR, albuminuria, hyperkalemia, volume status, or medication dose changes may affect monitoring and safety.

4. Reassess the regimen, not just the HbA1c

When one outcome improves and another remains suboptimal, review whether the regimen still matches the patient's priorities, tolerability, and monitoring capacity rather than automatically intensifying glucose lowering.

CLINICIAN DISAGREEMENT

Complex diabetes care is becoming a problem of prioritization, sequencing, and implementation - not simply escalation.

The survey shows that HCPs' primary challenge is not finding information. But it is often made a harder clinical decision. Obesity is the comorbidity most often selected as complicating diabetes management, HbA1c still anchors most treatment decisions, and the most selected next step is adding an SGLT2 inhibitor. Yet the most selected reason to continue therapy despite suboptimal HbA1c remains cardiovascular risk reduction.

That is a meaningful shift in how treatment success is framed. It therapy may be viewed by cardiovascular, kidney, or weight-benefit profile another part of the treatment plan is not working as intended, even if glucose control looks acceptable in isolation. If the patient cannot get access to it, it cannot be used for long.

The most important clinical insight may be the gap between preference and practice. Respondents place better cardiovascular risk reduction, kidney protection, and weight reduction ahead of glycemic control. But they also identify poor adherence, affordability, access, and fragmented care as the biggest barriers to realizing those priorities. That suggests that the challenge is not always choosing the right drug. Sometimes it is choosing the drug that can actually be delivered, supported, and sustained.

The future of complex diabetes care will therefore depend on more than new therapies. It will depend on how well clinicians can match the right patient to the right treatment sequence, at the right time, with the right level of support. The survey suggests that HCPs already know what matters most. The next challenge is building systems that make those priorities easier to act on.

Keep glycemia in context	**Let comorbidity change the sequence**
Use HbA1c as a starting point, but integrate kidney, cardiovascular, weight, and safety priorities.	Let obesity, kidney disease, heart failure, and cardiovascular risk shape the "next" decision, not only HbA1c.
Define benefit beyond one metric	**Treat feasibility as part of efficacy**
Ask what outcome each therapy is meant to protect, and when glycemic control falls short.	Consider access, cost, tolerability, and adherence before assuming a plan will work.

Help future diabetes-care insights reflect the trade-offs HCPs manage in real practice.

Join the conversation to share professional perspectives on treatment sequencing, adherence, access, and long-term outcomes.

Join the Forum

REFERENCES AND DATA SOURCE

1. American Diabetes Association Professional Practice Committee. Introduction and Methodology: Standards of Care in Diabetes—2026. *Diabetes Care*. 2026;49(Supplement 1):S1–S8. Doi.org/10.2337/dc26-SINT
2. American Diabetes Association Professional Practice Committee. Improving Care and Promoting Health in Populations: Standards of Care in Diabetes—2026. *Diabetes Care*. 2026;49(Supplement 1):S1–S8. Doi.org/10.2337/dc26-SPOP
3. American Diabetes Association Professional Practice Committee. Obesity and Weight Management for the Prevention and Treatment of Diabetes: Standards of Care in Diabetes—2026. *Diabetes Care*. 2026;49(Supplement 1):S1–S8. Doi.org/10.2337/dc26-SOBES
4. American Diabetes Association Professional Practice Committee. Pharmacologic Approaches to Glycemic Treatment: Standards of Care in Diabetes—2026. *Diabetes Care*. 2026;49(Supplement 1):S1–S8. Doi.org/10.2337/dc26-SPHAR
5. American Diabetes Association Professional Practice Committee. Cardiovascular Disease and Risk Management: Standards of Care in Diabetes—2026. *Diabetes Care*. 2026;49(Supplement 1):S1–S8. Doi.org/10.2337/dc26-SCVD
6. American Diabetes Association Professional Practice Committee. Chronic Kidney Disease and Risk Management: Standards of Care in Diabetes—2026. *Diabetes Care*. 2026;49(Supplement 1):S1–S8. Doi.org/10.2337/dc26-SCKD

MASTLocate Global Healthcare Insights is a series of reports and articles published by MASTLocate, a division of Mosaic Research, LLC, dedicated to providing insights into healthcare market trends and professional opinions. Information and analysis are based on data collected from healthcare professionals and publicly available sources. MASTLocate does not endorse any products or services mentioned in this report.

The content of this report is protected by copyright. Unauthorized reproduction, distribution, or use of any portion of this report without prior written permission is prohibited. For permissions, contact info@mastlocate.com.