Dry Eye Disease Management: When Symptoms Persist Beyond First-Line Treatment

Ophthalmology perspectives on DED phenotype, diagnostic weighting, first-line treatment, persistent symptoms, adherence, real-world outcomes evaluation, inadequate global stabilization, and the novel crowd-more-permanent-pain.

PARAMETRIC · ENGAGING · FIRST-LINE THERAPY · EVALUATION · LONG-TERM CONTROL

61	6	83.6%	9006491
Survey Answers	Questions Asked	Completion Rate	Staff

Survey Answers: 61 ophthalmologists in Germany, Canada, Switzerland, Japanese DOI, UK · Canada · Italy · France · Germany. Substantive DED symptoms were answered by 61 ophthalmologists. Incremental, experienced, experienced, after specialists, and therapist fields were excluded from analysis.

BUT THIS BEFORE!

Dry eye treatment becomes difficult when persistent symptoms expose more than one active disease driver.

Dry eye disease often looks simple at first glance. Symptoms are common, and many patients can be managed with standard first-line therapy. But when symptoms persist despite treatment, the problem is rarely just dryness. It may be inflammation, tear film instability, ocular surface damage, neuropathic pain, or a combination of these factors.

Clinicians then have to decide whether the first-line plan was insufficient, whether adherence or affordability limited exposure to treatment, whether the original phenotype was misclassified, or whether a second-line therapy is justified but poorly sequenced.

The MDPA/Cara survey explores how ophthalmologists (N=61) make these decisions, after diagnosis and treatment initiation, which findings carry diagnostic weight, what they do when patients remain symptomatic, and how they evaluate whether a therapy is working well enough to maintain DED care over the real-world journey.

Survey Answers: 61 ophthalmologists in Germany, Canada, Switzerland, Japanese DOI, UK · Canada · Italy · France · Germany. Substantive DED symptoms were answered by 61 ophthalmologists. Incremental, experienced, experienced, after specialists, and therapist fields were excluded from analysis.

CLINICAL SURVEY

TEDS DEWS III reinforces a driver-based model of diagnosis and management rather than a single dry-eye pathway.

TEDS DEWS III, published in 2025, updates dry eye disease as a heterogeneous, multifactorial disease involving loss of tear film and/or ocular surface homeostasis, with symptoms and signs that may vary over time and can be driven by different mechanisms in different patients.

Its diagnostic methodology combines symptoms with objective evidence of tear hyperosmolarity and tear-core subclassification to identify etiologic subtypes. Its management framework emphasizes a stepwise approach, with first-line therapy, reassessment, and escalation to more targeted therapies when the symptoms are directed toward what is actually causing disease in the individual patient.

The framework is most valuable when clinicians treat dry eye disease as a problem to be diagnosed, not a symptom to be suppressed. Its key limitation is reason to reassess mechanism, implementation, and disease burden rather than simply move to a stronger version of the same therapy.

TEDS DEWS III, published in 2025, updates dry eye disease as a heterogeneous, multifactorial disease involving loss of tear film and/or ocular surface homeostasis, with symptoms and signs that may vary over time and can be driven by different mechanisms in different patients.

DIAGNOSIS BARRIERS

Six findings that show where dry eye management shifts from first-line support to phenotype-driven escalation.

58.5%
Mixed mechanisms dominate
Mixed mechanisms (ADE) is the phenotype most often selected as driving management.

47.2%
Staining carries diagnostic weight
Corneal staining is the single finding most often selected as influencing DED diagnosis.

60.4%
Preservative-free tears want care
Preservative-free artificial tears remain the most common initial treatment.

49.1%
Adherence explains early failure
Poor treatment adherence is the most frequently selected answer when persistent symptoms after first-line therapy.

67.9%
Symptoms drive continuation
Sustained symptom relief is the dominant factor cited for continuing DED therapy.

45.3%
Personalized algorithms lead the future
Personalized DED treatment algorithms are the development most often expected to transform care.

01

TARGETING

Mixed-mechanism disease is the dominant management reality

58.5% say mixed mechanism dry eye most often drives management.

CONTEXT

A dry eye diagnosis does not automatically identify the clinician's target. Many patients have symptoms that are driven by instability, inflammation, neuropathic pain, or a combination of these findings. Clinicians must therefore ask which mechanism is driving the patient's symptoms and which treatment should be prioritized.

WHAT THE DATA SHOWS

Mixed mechanisms (ADE) was the most common phenotype selected, at 58.5%. Tear hyperosmolarity was the most frequently selected single mechanism, at 19.8%. Neuropathic pain was selected by 11.3%, and inflammation by 10.4%.

THE INSIGHT

Phenotyping has to remain dynamic. A patient who begins as "dryness" may later present with inflammation, pain, or mixed features. Management must change as the disease evolves.

WHY IT MATTERS

A fixed phenotype keeps sequencing and combination more important than escalation by intensity alone.

02

DIAGNOSTIC WEIGHT

Ocular-surface staining is the finding clinicians most often use to anchor the diagnosis

47.2% select ocular-surface staining as the most influential diagnostic finding.

CONTEXT

DED is diagnosed from a combination of symptoms and objective evidence of tear film instability. Staining is one of the most visible signs of ocular-surface damage, but its meaning depends on the tear film and ocular-surface examination.

WHAT THE DATA SHOWS

The MDPA was identified by 47.2% of respondents as the single most influential diagnostic finding. Tear film breakup time followed at 19.8%, while tear hyperosmolarity and meibomian gland dropout were each selected by 16.4%.

THE INSIGHT

TFOS DEWS III recommends a standardized diagnostic approach and emphasizes that no single test is sufficient. The survey suggests clinicians remain relevant when symptoms and staining do not fully align.

WHY IT MATTERS

A diagnosis that is more useful when it changes the next treatment decision, not simply when it confirms that the ocular surface is abnormal.

Finding that most influences DED diagnosis

Ocular surface staining	47.2%
Tear film breakup time (TBUT)	19.8%
Meibomian gland dropout	16.4%
Schirmer test	16.4%

03

FIRST-LINE TREATMENT

Preservative-free artificial tears remain the usual starting point, even in a mixed-mechanism population

60.4% initiate preservative-free artificial tears first.

CONTEXT

First-line management is the basic, accessible, and broadly applicable, which makes it essential that any survey-wide DED presentation. But it is also where the most patients are managed.

WHAT THE DATA SHOWS

Preservative-free artificial tears were most frequently described as the initial treatment, at 60.4%. Lipid-containing artificial tears were selected by 22.6%, and preservative-containing artificial tears by 17.0%.

THE INSIGHT

The open comments add real-world nuance of treatment escalation, long-standing multimodal treatment, and patients whose symptoms remain despite standard therapy.

WHY IT MATTERS

Preservative-free options confirm the treatment was usually selected, affordable, tolerated, and matched to the active driver.

Treatment initiated first

Preservative-free artificial tears	60.4%
Lipid-containing artificial tears	22.6%
Preservative-containing artificial tears	17.0%

04

EARLY TREATMENT FAILURE

Poor adherence and persistent inflammation account for most perceived first-line failures

49.1% identify poor treatment adherence as the most common explanation for initial treatment failure.

CONTEXT

Treatment failure can be demanding—multiple daily doses, delayed onset of benefit, discomfort or irritation. At once, device-based procedures. Often not act out of complete with routine life. Incomplete response to therapy can also reflect persistent inflammation.

WHAT THE DATA SHOWS

At the same time, the most common presence of persistent inflammation, at 49.1%. Poor adherence was selected by 45.3%, while early discomfort was selected by 41.5%, and a general dissatisfied plan was not selected by 34.0%.

THE INSIGHT

The open comments add real examples of treatment escalation, long-standing multimodal treatment, and patients whose symptoms remain despite standard therapy.

WHY IT MATTERS

Poor adherence, confirm the treatment was usually selected, affordable, tolerated, and matched to the active driver.

What most commonly explains initial treatment failure

49.1%	45.3%	41.5%
Persistent inflammation	Poor adherence	Early discomfort

05

FREQUENT SYMPTOMS

When symptoms persist, clinicians split between suppressing inflammation and intensifying MGD treatment

32.1% escalate anti-inflammatory therapy and 32.1% intensify MGD-directed treatment.

CONTEXT

Persistent symptoms are a fork in the pathway. One route treats inflammation more aggressively, another targets eyelid and gland dysfunction. A third questions the original diagnosis or exposure pattern. It is worth focus to determine how to proceed.

WHAT THE DATA SHOWS

Anti-inflammatory escalation and MGD-directed refinement were the most frequently chosen next steps in the survey. It also highlights why persistent DED is not a single escalation ladder.

THE INSIGHT

Responsiveness becomes particularly important when symptoms are disproportionate to visible signs. Clinicians may need to look beyond the ocular surface for explanation, such as central sensitization, systemic contributors, or other factors that change the ocular-surface environment.

WHY IT MATTERS

Persistent symptoms may be the suspected driver, not merely the fact that symptoms are still present.

Primary next step for persistent DED symptoms

Escalate anti-inflammatory therapy	32.1%
Intensify MGD-directed treatment	32.1%
Reassess diagnosis and exposure pattern	17.0%
Switch to preservative-free artificial tears	11.3%
Refer for specialty care	7.5%

06

DIAGNOSTIC DARKROOMS

Refractory MGD and autoimmune-associated dry eye most often force a different treatment strategy

37.7% select refractory meibomian gland dysfunction as the subgroup that most changes treatment strategy.

CONTEXT

These subgroups change more than the prescription choice. They can alter visit frequency, testing, and the expectations for response, and how clinicians define success.

WHAT THE DATA SHOWS

Refractory MGD and autoimmune-associated dysfunction (SJS/KC, Sicca) ranked highest, at 37.7% and 32.1%, respectively. Neuropathic pain was selected by 20.8%, and ocular surface disease, 9.4%.

THE INSIGHT

Some cases require only slight adjustments to the standard plan. Others require a different treatment strategy altogether. Clinicians may need to switch from a focus on surface-based strategies, control tactics, or combination care.

WHY IT MATTERS

Refractory DED is often a diagnostic problem, not a treatment problem. Not all access problem at the same time.

Subgroup that most changes treatment strategy

Refractory MGD	37.7%
Autoimmune-associated dysfunction (SJS/KC, Sicca)	32.1%
Neuropathic pain	20.8%
Ocular surface disease	9.4%

07

LONG-TERM CONTROL

Sustained symptom improvement matters more than an isolated surface metric when deciding whether to continue treatment

61.5% consider sustained symptom improvement as the moment that most supports long-term therapy.

CONTEXT

DED is a symptomatic disease, and patients judge treatment through comfort, visual function, functional burden, and day-to-day relief. Objective findings matter because they can help explain why symptoms are present.

WHAT THE DATA SHOWS

The survey most strongly supports a treatment that improves lasting symptoms. Sustained symptom improvement was selected by 61.5%, while symptom improvement without full normalization of signs by still required confirmation if the ocular surface is the dominant problem.

THE INSIGHT

Open-comment cases describe both directions of movement: patients whose surface improved but symptoms persisted, and patients whose comfort improved despite limited staining or inflammation.

WHY IT MATTERS

Clinicians may need to watch patients for symptoms, function, ocular surface signs, safety, and the specific mechanism being treated.

Outcome that most supports continuing long-term treatment

61.5%	23.1%	15.4%
Sustained symptom improvement	Symptom improvement without full normalization of signs	Improvement in one ocular surface measure

08

ADHERENCE CARE BARRIERS

Adherence and affordability remain the practical ceiling on advanced DED management

49.1% select poor patient adherence as the biggest barrier to advanced DED management.

CONTEXT

Advanced DED care can involve multiple prescriptions, procedures, repeated visits, dry therapy, ongoing symptoms, and long-term monitoring. Even when the treatment burden grows at the same time that symptoms may be the most disruptive.

WHAT THE DATA SHOWS

Poor adherence and administrative barriers outweigh technology barriers. Cost, access, and affordability can determine whether the plan is to be tried, adherence and affordability can determine whether the plan is actually followed.

THE INSIGHT

Open-comment examples include shared examples of insurance and coverage, and patients who can't afford the recommended drops or procedures. Patients should always underlying or chronic disease.

WHY IT MATTERS

Adverse and adherence are not administrative footnotes. They are variables that can alter the clinical outcome and should be addressed as part of treatment selection.

Biggest barrier to advanced DED management

Poor patient adherence	49.1%
Financial difficulties	37.7%
Limited access to procedures	9.4%
Limited access to procedures	3.8%

09

WHAT COMES NEXT

Personalized treatment algorithms lead the future vision for dry eye care

45.3% select personalized DED treatment algorithms as the development most likely to transform care.

Development most likely to transform DED care within 3–5 years

45.3%	28.3%	15.1%	11.3%
Personalized DED treatment algorithms	Novel anti-inflammatory therapies	Regenerative medicine approaches	Digital monitoring tools

CONTEXT

The preference for personalization for the rest of the survey. Mixed mechanisms dominate, symptoms drive continuation, and adherence and affordability are the biggest barriers to advanced care. Different clinicians may have different definitions of what personalization means, and different opinions about which tools, symptoms, and responses, and patient characteristics should guide it.

WHAT THE DATA SHOWS

Personalized DED treatment algorithms were the most frequently selected development likely to transform care, at 45.3%. Novel anti-inflammatory therapies were selected by 28.3%, while regenerative medicine approaches and digital monitoring tools were each selected by 13.2%.

THE INSIGHT

The survey does not suggest that algorithms should replace clinical judgment. It suggests a need for simpler, multimodal DED decisions more consistent and individualized.

WHY IT MATTERS

The future value of personalization will be measured by whether it reduces trial-and-error while keeping treatment practical for the patient.

OPEN-RESPONSE CASE

Real cases show that persistent dry eye often becomes a search for the driver that the first treatment did not address

Respondents were asked to describe challenging DED cases with persistent symptoms despite treatment, and to explain what mechanism, examination, and care decisions helped them identify the underlying problem.

Refractory disease remained difficult even after treatment
Several cases described patients who remained symptomatic despite multiple therapies, including punctal occlusion, autologous serum, and scleral lenses. Persistent pain, photophobia, or blurry vision changed the way clinicians framed the problem and the treatment approach.

Inflammation and systemic disease changed the ocular-surface picture
Examples included Sjögren syndrome, ocular rosacea, blepharitis, and ocular graft-versus-host disease. Clinicians described how inflammation, lid disease, and systemic contributors can change the clinical picture and require a different treatment strategy.

Symptoms and signs did not always improve together
Some patients had persistent symptoms despite apparently adequate tear production or ocular-surface staining. Others had improved signs but continued to report pain, dryness, or visual fluctuation. These cases pushed clinicians to look beyond the ocular surface for explanation, including central sensitization, systemic contributors, and neuropathic pain.

Neuropathic symptoms required a different pathway
Several cases described neuropathic pain, central sensitization, and symptoms that did not match the visible ocular surface. Clinicians described referral to pain clinics, neurology, and ophthalmology as part of the diagnostic pathway.

Access and insurance restrictions changed treatment
Some cases were shaped by insurance coverage, specialist availability, and limited access to advanced therapies. Clinicians described how patients had to switch from one treatment to another or adapt their care because of cost or coverage.

Recommended examinations revealed a different driver
Clinicians described examinations that changed the diagnosis, including meibography, tear osmolarity, tear breakup time, and corneal sensation. These findings helped explain why standard therapy had failed and guided more targeted treatment.

PEOPLE ALSO ASK · TOPIC FAQ

Frequently asked questions about dry eye disease diagnosis and management.

What is dry eye disease and how is it diagnosed?
Dry eye disease is a multifactorial disease of the tears and ocular surface that can cause symptoms of discomfort, visual disturbance, and tear film instability. It is usually diagnosed using a combination of symptoms and clinical signs.

What is mixed-mechanism dry eye?
Mixed-mechanism dry eye describes patients in whom more than one etiology drives symptoms or disease, such as tear hyperosmolarity, inflammation, eyelid abnormalities, or neuropathic pain. Management is therefore usually polytherapy, and when clinicians select their most dominant mechanism.

How is meibomian gland dysfunction related to dry eye disease?
Meibomian gland dysfunction is a common cause of evaporative dry eye. It can be assessed by meibography, tear osmolarity, tear breakup time, and corneal sensation.

What is anti-inflammatory treatment considered in dry eye disease?
Anti-inflammatory therapy is often considered when inflammation is an important disease driver or when symptoms persist despite lower-level support treatment. Choice of treatment depends on the clinical phenotype, severity, contraindications, access, and response to prior therapy.

Why may dry eye symptoms persist despite treatment?
Persistent symptoms can reflect incomplete adherence, ongoing inflammation, undiagnosed meibomian gland dysfunction, environmental or medication-related triggers, or pain mechanisms that are not fully explained by tear film or ocular surface findings alone.

What is neuropathic ocular pain in dry eye disease?
Neuropathic ocular pain is pain that may persist despite improvement in corneal disease and may be characterized as a central sensitization pain. It requires a different diagnostic approach, including careful history taking and exclusion of other causes.

ARTICLE SELECTION · PRACTICE CASE

What this means at four decision points in the dry-eye consultation.

The survey suggests that DED management succeeds when clinicians treat dry eye as a diagnostic problem, not a symptom to be suppressed. Four practical recommendations, in a consultation, ophthalmologist or optometrist.

1. Reconfirm the active drivers.
Measure tear osmolarity, assess the lid anatomy, MGD, ocular-surface staining, and corneal sensitivity before choosing therapy. Mixed mechanisms is the most common phenotype in this survey.

2. Separate nonresponse from nonuse.
Ask whether symptoms persist despite adherence, or whether adherence, cost, and access barriers are preventing the therapy from working.

3. Track symptoms and signs in parallel.
Review the ocular-surface exam, symptom burden, and treatment burden together. Dry eye can improve on one measure while remaining poor on another.

4. Recognize refractory subgroups early.
Refractory MGD, autoimmune-associated dysfunction (SJS/KC, Sicca), and neuropathic pain may require a different strategy than standard first-line therapy.

GLOBAL PERSPECTIVE

The hardest dry-eye cases are rarely asking for "more treatment"; they are asking for a better match between mechanism, burden, and what the patient can sustain.

The survey confirms that ophthalmologists diagnose dry eye as a multifactorial disease driven by tear film instability, inflammation, and ocular-surface dysfunction. Preservative-free tears remain the common first step, but when symptoms persist, the pathway fractures quickly into inflammation, meibomian gland dysfunction, neuropathic pain, and adherence barriers.

The strongest theme is not quantity of treatment, but mismatch. Clinicians are not failing because they lack options. They are struggling because persistent symptoms can have more than one cause, and because the first treatment rarely addresses all of them. Mixed mechanisms (ADE) was the most frequently selected phenotype, while tear hyperosmolarity, inflammation, and neuropathic pain each remained important. Treatment must therefore be sequenced, not simply escalated.

The data also highlight the importance of chronicity, progression, and symptom-sign mismatch. Inadequate response, adverse experience (AE/SAE), neuropathic pain, and access barriers can all change what "success" means and how a patient may need individualized care.

Finally, the survey suggests that personalized DED treatment algorithms are the future development expected to transform care. In real-world care, it is the possibility of matching the right test to the symptoms, phenotype, diagnostic findings, symptom burden, and treatment goals. Personalization is not about more treatment. It is about fewer wrong turns.

Phenotype-driven DED management is the future.
Personalized DED care is expected to transform how dry eye disease is managed.

Symptoms and signs together
Symptoms and signs must be considered together.

Escalate according to the driver, not the next crown
Escalate according to the driver, not the next crown.

Help future ocular-surface insights reflect the complexity of real-world dry eye care.
Join clinicians in three clinical perspectives on DED diagnosis, phenotype, treatment response, maintenance goals preferences, adherence, and real-world outcomes.
Join the Panel

REFERENCES AND DATA SOURCE

1. **MDPA/Cara Ophthalmology Survey.** Dry Eye Disease Management: When Symptoms Persist Beyond First-Line Treatment. Conducted September 2025. Survey Answers: 61 ophthalmologists in Germany, Canada, Switzerland, Japanese DOI, UK · Canada · Italy · France · Germany. Substantive DED symptoms were answered by 61 ophthalmologists. Incremental, experienced, experienced, after specialists, and therapist fields were excluded from analysis.
2. **TFOS DEWS III.** The Ocular Surface. 2025. Tear Film & Ocular Surface Society. DOI: Science.
3. **TFOS DEWS III Executive Summary.** Ocular Surface. 2025 of Ophthalmology. 2025. Science.

MDPA/Cara Ophthalmology Survey. Dry Eye Disease Management: When Symptoms Persist Beyond First-Line Treatment. Conducted September 2025. Survey Answers: 61 ophthalmologists in Germany, Canada, Switzerland, Japanese DOI, UK · Canada · Italy · France · Germany. Substantive DED symptoms were answered by 61 ophthalmologists. Incremental, experienced, experienced, after specialists, and therapist fields were excluded from analysis.

Powered by Cara Ophthalmology · Privacy Policy · Terms of Use · Contact Us