Epilepsy in Pregnancy: Balancing Seizure Control and Medication Safety

A cross-market view of how neurology HCPs approach pregnancy planning, antiseizure medication selection, fetal safety, therapeutic monitoring, seizure recurrence, and postpartum dose management.

Preconception planning · ASM selection · Pregnancy monitoring · Dose adjustment · Postpartum transition

112	6	61.6%	9010428
Survey respondents	Countries fielded	Completion rate	Fielded

Japan, UK, Canada, Italy, France, Germany

Illustrative example. Fictional company, country names, and data. Not for distribution.

WHAT THIS REPORT

Epilepsy management in pregnancy is a sequence of decisions, not a single medication choice.

For HCPs, the neurology-OB/GYN conversation can begin years before pregnancy and can change again at conception, during each trimester, and after delivery. The challenge is to balance seizure control, medication exposure, and postpartum monitoring across connected visits—while recognizing that pregnancy itself can change how epilepsy behaves.

That makes timing central. A planned pregnancy creates space to review medication, reduce exposure, time folic acid, reassess seizure risk, and prepare for monitoring. An unplanned pregnancy often forces those decisions into a period when fetal exposure may already be underway and when pregnancy changes may carry its own risk.

The REPLICAS survey explores where physicians see the greatest challenges in managing epilepsy management, what drives preconception medication decisions, how pregnancy affects monitoring, and what unquestions believe could make care safer over the next several years.

METHODOLOGY

A cross-market survey of professional neurology and epilepsy-care HCPs.

This study included professional HCP respondents across the US, UK, Italy, France, and Germany. The source file contained 112 survey records with a 61.6% completion rate. Question-level response bases varied as respondents progressed through the survey.

The response base is the numerical source of each percentage. Excluded and non-applicable responses are noted where relevant. Multi-select questions are identified on the relevant slides. Percentages may not sum to 100% due to rounding.

EXECUTIVE SUMMARY

Six signals define the reproductive epilepsy-management challenge.

01

42.4% identify *unplanned pregnancy* as the leading challenge.

The biggest challenge is the management of patients who enter pregnancy without a single medication-safe effect.

02

71.6% say *planned pregnancy* prompts SOM review.

Planned pregnancy is where the strongest opportunity for reviewing the regimen.

03

Fetal safety dominates preconception selection.

Improved teratogenicity risk and fetal neurodevelopment are the most cited factors for ASM choice.

04

42.9% say *falling serum levels* most alter pregnancy management.

Drug clearance can change even when the prescribed dose does not.

05

49.6% want *safer pregnancy compatible ASMs*.

Respondents see medication exposure itself as the development most likely to transform care.

PATIENT JOURNEY

Six signals define the reproductive epilepsy-management challenge.

01

42.4% identify *unplanned pregnancy* as the leading challenge.

The biggest challenge is the management of patients who enter pregnancy without a single medication-safe effect.

02

71.6% say *planned pregnancy* prompts SOM review.

Planned pregnancy is where the strongest opportunity for reviewing the regimen.

03

Fetal safety dominates preconception selection.

Improved teratogenicity risk and fetal neurodevelopment are the most cited factors for ASM choice.

04

42.9% say *falling serum levels* most alter pregnancy management.

Drug clearance can change even when the prescribed dose does not.

05

49.6% want *safer pregnancy compatible ASMs*.

Respondents see medication exposure itself as the development most likely to transform care.

PATIENT JOURNEY

Pregnancy planning changes the epilepsy-management frame more than any other reproductive presentation.

56.3% select *pregnancy planning* as the reproductive presentation that most often changes management.

CONTEXT

A patient can be seriously stable for years and still require a different management approach when pregnancy is planned. Preconception counseling, contraception, folic acid, seizure history, and the feasibility of changing therapy all become part of the same decision.

WHAT THIS DATA SHOWS

Pregnancy planning leads at 56.3%, followed by breakthrough seizures at 23.8%, unplanned pregnancy at 10.2%, and contraception planning at 9.7%.

THE RESULT

The result suggests the reproductive intent is not merely contextual information. It changes how we interview stable epilepsy patients.

WHY IT MATTERS

The most recommended guideline emphasizes optimizing antiseizure medication before conception and continuing treatment during pregnancy. The survey finding reflects the practical reason: the visit is the moment a regimen is often tested against pregnancy considerations.

Reproductive presentation that most changes epilepsy management

Pregnancy planning	56.3%
Breakthrough seizures	23.8%
Unplanned pregnancy	10.2%
Contraception planning	9.7%

WHEN THE REGIMEN GETS REVIEWED

Planned pregnancy is the strongest trigger for antiseizure medication review.

71.6% select *planned pregnancy* as a factor prompting SOM review.

CONTEXT

Medication review is anything but often triggered by loss of seizure control or adverse effects. Pregnancy planning adds another trigger because the risk-benefit profile of the same medication can change when fetal exposure becomes relevant.

WHAT THIS DATA SHOWS

Planned pregnancy leads at 71.6%, followed by *breakthrough seizures* at 61.3%, *contraception interactions* at 55.2%, and *adverse effects* at 48.7%.

THE RESULT

Contraception interactions are neither because some enzyme-inducing ASMs can reduce the effectiveness of hormonal contraceptives. Pregnancy planning therefore makes a more pregnancy-centered, medication-focused conversation.

WHY IT MATTERS

Contraception interactions also matter because some enzyme-inducing ASMs can reduce the effectiveness of hormonal contraceptives. Pregnancy planning therefore makes a more pregnancy-centered, medication-focused conversation.

Factors that prompt antiseizure medication review

Planned pregnancy	71.6%
Breakthrough seizures	61.3%
Contraception interactions	55.2%
Adverse effects	48.7%

PRECONCEPTION MEDICATION REVIEW

Preconception choice is dominated by fetal risk, with valproate and neurodevelopment at the center.

49.6% cite *improved teratogenicity risk* and 44.2% cite *improved fetal neurodevelopmental risk*.

CONTEXT

Preconception counseling is where risk is already the medication with the lowest documented fetal risk. It is the safest regimen that is also appropriate for the patient's seizure history, lifestyle, and reproductive goals.

WHAT THIS DATA SHOWS

Improved teratogenicity risk and *fetal neurodevelopmental risk* were chosen by 49.6% of respondents, followed by *seizure control efficacy* at 41.3% and *seizure control efficacy* by 38.9% in the single-select question.

THE RESULT

The forecast choice format makes that safety especially visible, but it should not be interpreted as clinicians ignoring seizure control. Seizure control is the baseline requirement; teratogenicity and neurodevelopment are the factors that most often create a decision.

WHY IT MATTERS

Current guidance reinforces that patients should not stop ASMs without medical supervision, but it also underscores that unplanned pregnancy therefore needs to prioritize the lowest-risk regimen that maintains seizure control. Valproate, lamotrigine, and levetiracetam are the most relevant examples.

ASM factor that most influences preconception selection

Improved teratogenicity risk	49.6%
Improved fetal neurodevelopmental risk	44.2%
Seizure control efficacy	41.3%
Seizure control efficacy	38.9%

TREATMENT TRANSITION

Unplanned pregnancy removes the time clinicians need to optimize reproductive epilepsy care.

49.4% identify *unplanned pregnancy* as the factor that most commonly complicates management.

CONTEXT

Preconception planning works only when pregnancy intention is known early enough to act on it. Unplanned pregnancy is the scenario in which that window has already closed. Folic acid may not have started, and the patient may be simultaneously anxious about continuing treatment.

WHAT THIS DATA SHOWS

Unplanned pregnancy leads at 49.4%, followed by *ASM adherence* at 27.4%, *breakthrough seizures* at 13.3%, and *inadequate folic acid use* at 9.9%.

THE RESULT

The finding underscores a timing failure. It turns decisions that could have been addressed into decisions that must be made under time pressure after pregnancy has already started.

WHY IT MATTERS

Current guidance emphasizes that patients should not stop ASMs without medical supervision, but it also underscores that unplanned pregnancy therefore needs to prioritize the lowest-risk regimen that maintains seizure control. Valproate, lamotrigine, and levetiracetam are the most relevant examples.

What most commonly complicates reproductive epilepsy management

Unplanned pregnancy	49.4%
ASM adherence	27.4%
Breakthrough seizures	13.3%
Inadequate folic acid use	9.9%

TREATMENT TRANSITION

Switching to a lower-risk ASM leads, but one-size-fits-all switching does not.

55.8% select *switching to a lower-risk ASM* as the primary strategy before pregnancy.

CONTEXT

Preconception medication changes are most useful when there is enough time to taper, titrate, and monitor. Switching can reduce exposure, but it can also destabilize seizure control if it is rushed or poorly timed.

WHAT THIS DATA SHOWS

Switching to a lower-risk ASM leads at 55.8%, followed by *add adjunctive ASM* at 31.2%, *maintain current regimen* at 9.7%, and *taper to monotherapy* at 3.3%.

THE RESULT

The spread across the three options suggests that clinicians see the trade-off against by 55.7%, *maintaining effective monotherapy* by 31.2%, and *optimizing folic acid timing* by 13.1%.

WHY IT MATTERS

The clinical priority is the lowest risk regimen that still controls the patient's seizures. That is why the same medication can be appropriate for one patient and contra-hazardous choice for another.

Primary ASM strategy before pregnancy

Switch to lower-risk ASM	55.8%
Add adjunctive ASM	31.2%
Maintain current regimen	9.7%
Taper to monotherapy	3.3%

TREATMENT TRANSITION

Falling serum ASM levels can become the first sign that a stable regimen is no longer stable.

42.9% say *falling serum ASM levels* most alter management during pregnancy.

CONTEXT

Pregnancy can alter clearance and distribution of antiseizure medications. For some patients, that means a dose that was working well can quietly stop working. Knowing the risk that a patient who was previously controlled begins to experience breakthrough seizures.

WHAT THIS DATA SHOWS

Falling serum ASM levels leads at 42.9%, breakthrough *generalized seizures* at 31.7%, *breakthrough focal seizures* at 17.7%, and *hyperemesis affecting absorption* at 7.7%.

THE RESULT

The survey makes therapeutic monitoring clinically concrete. The answers to this question suggest that clinicians are not just watching for seizures. They are watching for pharmacokinetic changes that can precede them.

WHY IT MATTERS

ACOG guidance recommends considering therapeutic drug monitoring concentrations and monitoring seizures before ASMs during pregnancy when clinically indicated. Falling levels are the kind of change that can be missed between appointments, unforeseen symptoms, and the patient's individual therapeutic baseline.

Pregnancy change that most alters ASM management

Falling serum ASM levels	42.9%
Breakthrough generalized seizures	31.7%
Breakthrough focal seizures	17.7%
Hyperemesis affecting absorption	7.7%

OTHER DEVELOPMENT MEDICATIONS AND CARE

Seizure recurrence, fetal safety, and serum levels all compete for attention during pregnancy.

52.3% select *seizure recurrence*, 50.0% *fetal safety*, and 38.6% *serum ASM concentrations* as adjustment factors.

CONTEXT

Pregnancy dose adjustment is rarely justified by one data point. A lower serum concentration may not matter if seizures remain stable, while dropping levels can be the earliest warning that a medication-pregnancy interaction needs to be addressed.

WHAT THIS DATA SHOWS

Seizure recurrence leads at 52.3%, *fetal safety* at 50.0%, and *serum concentrations* at 38.6%. Multiple factors can be selected.

THE RESULT

The leading factors describe a three-way clinical balance: what the patient is experiencing, what the medication level is doing, and what the fetus may be exposed to.

WHY IT MATTERS

This is why therapeutic drug monitoring is most useful when it is interpreted alongside seizure history, dose, and timing. It can help explain why the standard for planned is difficult to maintain while protecting unnecessary dose escalation.

Factors that guide ASM adjustment during pregnancy

Seizure recurrence	52.3%
Fetal safety concerns	50.0%
Serum ASM concentrations	38.6%
Hyperemesis affecting absorption	7.7%

POSTPARTUM TRANSITION

Delivery ends the pregnancy, not the medication-management problem.

53.3% favor *restoring the pre-pregnancy ASM dose and 28.9% favor checking postpartum serum levels*.

CONTEXT

If an ASM dose was increased during pregnancy to compensate for higher clearance, the postpartum period can be the moment when that dose becomes problematic as clearance returns toward baseline.

WHAT THIS DATA SHOWS

Restoring the pre-pregnancy dose and *checking postpartum serum levels* are tied at 53.3%. Adjusting according to seizure control is selected by 31.7%, and *maintaining dose* is selected by 15.8%.

THE RESULT

The split between dose restoration and level checking points to the same underlying issue: postpartum dose management is not just about seizure liberation, and passive continuation of the pregnancy regimen.

WHY IT MATTERS

Postpartum dose tapering can unmask what is shown medication-based pharmacokinetic changes that the dose after child-birth and changed during pregnancy. Clinical values, serum levels whereas, dose duration, and tissues all remain part of postpartum review.

Postpartum approach that best guides ASM management

Restore pre-pregnancy dose	53.3%
Check postpartum serum levels	28.9%
Adjust according to seizure control	31.7%
Continue pregnancy dose without change	15.8%

OTHER DEVELOPMENT MEDICATIONS AND CARE

Respondents want safer pregnancy-compatible ASMs more than another layer of technology.

49.6% select *safer pregnancy-compatible ASMs* as the development most likely to transform care.

CONTEXT

Digital tools can support preconception and monitoring, but safer ASMs can remove exposure, but they do not remove the fundamental tension between maintaining seizure control and minimizing fetal exposure.

WHAT THIS DATA SHOWS

Safer pregnancy-compatible ASMs leads at 49.6%, followed by *pregnancy registries* at 21.7%, *digital seizure prediction* at 17.6%, and *telemedicine follow-up* at 11.1%.

THE RESULT

The result reinforces a preference for a pharmaceutical shift rather than digital. Respondents appear to want more treatment options to reduce the need to choose between seizure stability and reproductive safety.

WHAT IT MEANS

Recent guidance and regulatory updates show how rapidly reproductive epilepsy is changing epilepsy prescribing. For example, the MHRA's decision that pregnancy prevention measures for topiramate in 2024. Future programs will need to track how quickly clinicians adopt new options as they become and how their access progresses.

Development that could most transform care within 3-5 years

Safer pregnancy-compatible ASMs	49.6%
Pregnancy registries	21.7%
Digital seizure prediction	17.6%
Drug monitoring	11.1%
Telemedicine follow-up	11.1%
Precision reproductive care	0.0%

OPEN-ENDED CODE LOOP

Real-world cases show how quickly reproductive epilepsy care becomes a balance between competing risks.

The open-ended question asked respondents to describe a challenging epilepsy case involving reproductive care, what changed management, and what remained unresolved.

Unplanned pregnancy: unapproved Brivaracetam dose

Woman stable, transitioned, decided pregnancy in her thigh after she was planning to conceive. Brivaracetam dose was increased during pregnancy. Unplanned pregnancy, dose was increased during pregnancy.

Medication switching could create new instability

Patient was stable on her medication but wanted to become pregnant. Medication switching always to create instability.

Fear of fetal harm could affect adherence

Patient was stable on her medication but was concerned about fetal harm. Fear of fetal harm could affect adherence.

Seizure control maximized the counterweight to fetal safety

Reproductive sexuality discreet difficulty. Changing medication in patients with epilepsy can be difficult. Seizure control maximized the counterweight to fetal safety.

Drug-food monitoring became more important during pregnancy

Patient was stable on her medication but was concerned about fetal harm. Drug-food monitoring with other issues started changed during pregnancy.

Multidisciplinary care helped when decisions were more complex

Patient was stable on her medication but was concerned about fetal harm. Multidisciplinary care helped when decisions were more complex. Also, to ensure ASM levels in the safe.

PATIENT JOURNEY

Common clinical questions about epilepsy in pregnancy.

Which antiseizure medications are considered when treating people with epilepsy who may become pregnant?

Lamotrigine and levetiracetam are the most commonly prescribed ASMs in pregnancy. Valproate is associated with substantial fetal harm and is not recommended for women of childbearing potential unless other medications have failed or no other options are available.

Why is valproate a major concern in pregnancy?

Valproate exposure in pregnancy is associated with major congenital malformations, neural tube defects, and impaired neurodevelopment in exposed children. Because of these risks, valproate is not recommended for women of childbearing potential unless other medications have failed or no other options are available.

Should antiseizure medication be stopped when pregnancy is confirmed?

In most cases, antiseizure medication should not be stopped. Seizures can be dangerous for the pregnant woman and fetus. Drug pregnancy has changes, changes in an effective regimen based on pregnancy-related changes.

Why can antiseizure medication levels fall during pregnancy?

Pregnancy can change drug absorption and distribution, lowering blood concentrations of some antiseizure medications even when the dose has not changed. For some medications, dose adjustments may be needed to maintain seizure control. Therapeutic drug monitoring can help guide these decisions.

Is folic acid recommended for people with epilepsy who may become pregnant?

Yes. Folic acid is recommended before conception and during pregnancy for people with epilepsy who may become pregnant. Some antiseizure medications can reduce folate levels, increasing the importance of supplementation.

How should antiseizure medication dosing be managed after delivery?

Patients should be counseled during pregnancy that medication clearance may increase soon after delivery. ACOG recommends seizures and antenatal care to review medications toward prescription dosages in the first few days after birth, with clinical review and serum level monitoring where appropriate.

PATIENT JOURNEY

What this means across the reproductive epilepsy-care pathway.

The survey suggests that the most important decisions are distributed across time. Good reproductive epilepsy care depends on anticipating the next transition, not just optimizing the medication at the current visit.

1. Ask about pregnancy intention before it becomes urgent

Regular reproductive-planning review visits to review ASM risk, contraception, and folic acid status. Pregnancy intention should be a routine part of epilepsy care.

2. Use monitoring to interpret clinical change

Therapeutic drug monitoring can help distinguish pharmacokinetic change from nonadherence. Serum levels are most useful when interpreted alongside seizure history.

3. Plan the postpartum dose transition before delivery

Postpartum dose tapering can unmask what is shown medication-based pharmacokinetic changes that the dose after child-birth and changed during pregnancy. Clinical values, serum levels whereas, dose duration, and tissues all remain part of postpartum review.

OPEN-ENDED CODE LOOP

The central challenge is not pregnancy alone. It is managing epilepsy across changing reproductive states without losing seizure control.

The REPLICAS challenge point is a care pathway in which timing matters as much as medication choice. Planned pregnancy opens a window for deliberate review. Unplanned pregnancy must begin with that window and focus on reference-setting fetal exposure against the risk of pregnancy and may be the presenting breakthrough seizures.

The survey also shows the pregnancy is managed at a date, not one-point event. Drug levels may fall, seizure recurrence may emerge, fetal safety concerns may intensify, and postpartum dose tapering may unmask changes that were invisible during pregnancy. The most useful clinical question is often not "what dose?" but "what has changed since the last visit?"

Postpartum care is the last transition, not the end of monitoring. Dose reductions should be tied to both breast-pharmacokinetic levels, and patients should be counseled that seizure risk can return quickly after delivery.

The strongest future preferences — safer pregnancy-compatible ASMs, capacity for unmanaged need behind the entire survey. Better monitoring and digital prediction tools are also important, but they are secondary to medications that reduce fetal exposure without increasing seizure risk. Pregnancy registries and telemedicine follow-up are also important, but they are further down the list.

Normalize preconception review

Make reproductive planning part of routine epilepsy visits, not a separate discussion. Ask about pregnancy intention, contraception, and folic acid status at regular intervals.

Individualize medication risk

Choose medications based on seizure history, pregnancy intention, and fetal exposure. Valproate, lamotrigine, and levetiracetam are the most relevant examples.

Monitor the patient, not the level

Interpret serum concentrations alongside seizure frequency, dose changes, and clinical symptoms. Therapeutic drug monitoring can help distinguish pharmacokinetic change from nonadherence.

Build the postpartum plan before delivery

Postpartum dose tapering can unmask what is shown medication-based pharmacokinetic changes that the dose after child-birth and changed during pregnancy.

Help future neurology insights reflect the reproductive-care decisions HCPs make every day.

Join our Panel

Share your perspectives and influence what we commission in the evidence-to-commercial shaping roadmaps.

REFERENCES AND DATA SOURCE

1. ACOG Practice Bulletin No. 208: Update on Treatment of Epilepsy During Pregnancy. Obstet Gynecol. 2019;133(5):e305-e319.
2. American Academy of Neurology, American Epilepsy Society. Practice Advisory: Updated Evidence-Based Guidance on Antiseizure Medication Use in Pregnant Women with Epilepsy. Neurology. 2020;94(9):e889-e900.
3. Medicines and Healthcare products Regulatory Agency. Valproate: new restrictions on use in women and girls. MHRA Drug Safety Update. 2018.
4. Medicines and Healthcare products Regulatory Agency. Topiramate: new safety measures including pregnancy prevention. 2024. MHRA topiramate safety update.