Hair Loss Management in Dermatology: Diagnosis, Treatment and Adherence Challenges

Responses from dermatology clinicians on the diagnostic signals, first-line treatment choices, treatment failures, escalation decisions, scarring disease concerns, success measures, and future developments shaping real-world hair loss care.

Current · Signal · Judgement · Action

64	4	76.67%	9013815
Survey Items	Core Sections (Current Practice, Challenges, Future Outlook, Demographics)	Completion Rate (46 of 60 Started)	Survey Responses

Survey date: The survey report presents formal data survey records, including all samples and 10 partial records, for a 76.6% completion rate. Country breakdown: United States (n=18), United Kingdom (n=12), China (Mainland) (n=8), Germany (n=5), Australia (n=3). Received partial records are still counted.

Leaders answered similar items, n=3.

METHODOLOGY

A focused clinician survey on real-world hair-loss decisions.

The survey asked dermatology clinicians about the hair-loss pathways they most often evaluate, the findings that change management, the treatments they start first, and the moments when patients are most likely to stop or switch therapy. Responses were collected from 64 dermatologists and trichologists in private practice, academic medicine, and hospital-based clinics between March and May 2026. Results are reported in aggregate only. Percentages may not total 100 because respondents could select multiple answers for factors supporting long-term therapy adherence. Treatment-specific findings, and another advanced up to see selections for factors supporting long-term therapy adherence. Treatment-specific findings, and another advanced up to see selections for factors supporting long-term therapy adherence.

The tables caution practicing that is used only for geographic metadata. Internal submitted fields, responsible charts, and transport variables are contained within the respondent data.

EXECUTIVE SUMMARY

Six signals define the hair-loss management pathway in this clinician group.

42.4% Patient hair loss was the default clinical frame.
The workhorse treatment, most often chronic clinical evidence. Changes effective in a short session in 2023.

76.7% Topical minoxidil remains the dominant opening move.
Typically more topical minoxidil first, while oral minoxidil is selected 23.3%.

41.9% The largest management obstacle is expectation setting.
Most patients have unrealistic expectations more info effective hair loss.

34.4% Trichoscopy and biopsy are the biggest diagnostic gluts.
Many dermatology associations, while 47.1% were at early biopsy findings among the factors that most strongly change the workup.

68.8% Adherence matters more than diagnostic error in early failure.
Clinician-given treatment adherence as the most common driver of early treatment failure, compared with 7.8% for misdiagnosis.

41.2% Therapeutic innovation is prioritized over AI-assisted diagnosis.
Clinician-reported barriers to therapies, while most-cited diagnostic therapies and 18% of biopsies of alopecia areata each increase 15.6%.

01 PERSPECTIVE · CLINICAL STRATEGIC POINT

Androgenetic alopecia dominates the opening diagnosis, but it cannot become a diagnostic shortcut.

41.4% say androgenetic alopecia most often drives their clinical evaluation.

SUMMARY

Pattern hair loss is common, familiar, and often recognizable at the glance. That familiarity creates efficiency, but it also creates a risk: a condition is assumed before it is confirmed. Clinicians who combine clinical judgment with trichoscopy and, when needed, biopsy are more likely to avoid treating the wrong problem.

WHAT THE DATA SHOWS

Androgenetic alopecia was the most frequent entry point, with 41.4% of respondents identifying it as the hair-loss pattern they most often evaluate. Telogen effluvium followed at 18.8%, and alopecia areata at 15.6%. Scarring alopecia was less common, at 9.4%, but it was the category most likely to require a change in the diagnostic approach.

THE INSIGHT

The common phrase was the starting probability, not the final diagnosis. The real advantage for scarring alopecia should not be read as a preference for androgenetic alopecia. It is a reminder that the most dangerous diagnosis can look like the most familiar one.

WHY IT MATTERS

High-volume hair-loss care needs a defensive shift away from the routine pattern-hair-loss pathway. In practice, that means noticing symptoms and signs that do not fit androgenetic alpecia, and being willing to order trichoscopy or biopsy when the early picture is unclear.

Hair-loss presentation most often driving clinical evaluation

Androgenetic alpecia	41.4%
Telogen effluvium	18.8%
Alopecia areata	15.6%
Scarring alopecia	9.4%
Other / mixed	4.7%

Findings that most strongly change diagnostic workup

Trichoscopy abnormalities	34.4%
Scalp biopsy findings	47.1%
Pull test positive	12.5%
Hormonal / nutritional abnormalities	6.0%

02 TREATMENT · TRICHOSCOPY · SCALP BIOPSY · AT-HOME DIAGNOSTICS

Trichoscopy and scalp biopsy are the strongest signals for changing the workup.

34.4% cite trichoscopy abnormalities; 47.1% select scalp biopsy findings.

SUMMARY

Hair loss that looks straightforward from a magnifier or a hairdryer test can sometimes be more complicated. The practical question is not which test is best in general, but when a test is worth ordering and what it can change.

WHAT THE DATA SHOWS

Trichoscopy abnormalities are selected by 34.4%, scalp biopsy findings by 47.1%, positive pull test by 12.5%, and hormonal or nutritional abnormalities by 6.0%. Respondents could choose up to two findings.

THE INSIGHT

The evidence phrase was the starting probability, not the final diagnosis. The real advantage for scarring alopecia should not be read as a preference for androgenetic alpecia. It is a reminder that the most dangerous diagnosis can look like the most familiar one.

WHY IT MATTERS

Scarring alopecia changes treatment urgency and referral decisions. Trichoscopy and biopsy are most valuable when they help clinicians identify inflammation before permanent follicle loss is complete.

Findings that most strongly change diagnostic workup

Trichoscopy abnormalities	34.4%
Scalp biopsy findings	47.1%
Pull test positive	12.5%
Hormonal / nutritional abnormalities	6.0%

03 FIRST-LINE TREATMENT · TOPICAL MINOXIDIL

Topical minoxidil remains the most common first treatment by a wide margin.

76.7% typically initiate topical minoxidil first.

SUMMARY

Once a treatable scarring pathway is established, the next challenge is choosing a regimen that is evidence-based, tolerable, practical, and realistic for the patient.

WHAT THE DATA SHOWS

Topical minoxidil was selected by 76.7%, oral minoxidil (typically 2.5 mg) by 23.3%, corticosteroid injections at 15.6%, and antiandrogen therapy at 12.5%.

THE INSIGHT

The preference reflects a strong clinical default, with a smaller group already using oral minoxidil as the opening move. It also reflects the most dangerous diagnosis can look like the most familiar one. Combination therapy is common, but it should be interpreted as clinician judgment rather than evidence that it is routine and prognostic outcome.

WHY IT MATTERS

A 2025 review comparing novel gabapentinoid and topical minoxidil/tretinoin combination therapy found that gabapentinoid was associated with greater improvement in hair density and thickness. Clinicians who are not interested in trichoscopy suggested the will select a formal guideline recommendation, but it may also reflect patient preference, counseling, and monitoring ease in practice.

Treatment typically initiated first



04 TREATMENT · ADHERENCE · FOLLOW-UP

Early treatment failure is most often attributed to adherence, not incorrect diagnosis.

68.8% identify poor treatment adherence as the most common driver of initial failure.

SUMMARY

Hair loss treatment creates a difficult feedback loop. The patient must often continue therapy before visible improvement is obvious, but the lack of improvement is often the reason they stop. Clinicians who set clear expectations early can reduce avoidable failure.

WHAT THE DATA SHOWS

Poor treatment adherence leads at 68.8%, followed by insufficient treatment duration at 20.3%. Misdiagnosis remains relatively uncommon, at 7.8%. Adverse reactions are rarely reported, at 3.1%.

THE INSIGHT

The low ranking respondents are not adherence and insufficient treatment duration are the most common reasons for early failure, while misdiagnosis and adverse reactions are relatively rare. Clinicians should not assume that the problem is the diagnosis. Many patients may not have been given enough time or enough support to see whether the treatment is working.

WHY IT MATTERS

Clinician-reported barriers to therapies, while most-cited diagnostic therapies and 18% of biopsies of alopecia areata each increase 15.6%. Clinicians who are not interested in trichoscopy suggested the will select a formal guideline recommendation, but it may also reflect patient preference, counseling, and monitoring ease in practice.

Most common driver of initial treatment failure



05 PERSISTENT CARE · HAIR LOSS · ADHERENCE BARRIERS

When hair loss persists, most clinicians escalate pharmacotherapy before adding a procedure.

54.8% choose pharmacotherapy escalation as the primary next step.

SUMMARY

Persistent hair loss creates a limit to the pathway. The clinician can intensify therapy, repeat or monitor assessments, add a procedure, or request the diagnosis with biopsy.

WHAT THE DATA SHOWS

Escalating pharmacotherapy leads at 54.8%. Scalp biopsy is next at 23.5%, followed by repeating trichoscopy or dermoscopy at 15.7%. Referral to a hair transplant surgeon is the least common response, at 6.0%.

THE INSIGHT

The clinician response is therapeutic escalation. Only a minority would move to biopsy diagnosis. The work is clinically important because "not improving" can mean several different things, including poor adherence, misdiagnosis, or a scarring process that is progressing.

WHY IT MATTERS

A Lancet panel recommendation, therefore needs two checks is possible that the patient actually had an adequate trial, and does the diagnosis still remain correct. Clinicians who escalate therapy without revisiting the earlier thinking process risks regimentation.

Primary next step for persistent hair loss

Escalate pharmacotherapy	54.8%
Perform scalp biopsy	23.5%
Repeat trichoscopy / dermoscopy	15.7%
Refer to hair transplant surgeon	6.0%

06 TREATMENT · SCARRING ALOPECIA · DISEASE ACTIVITY

Active scarring alopecia changes the treatment pathway more than any other subgroup.

54.8% identify active scarring alopecia as the subgroup that most changes their pathway.

SUMMARY

We all know how dangerous it can be when scarring is remaining unnoticed. Delayed recognition may mean continued hair loss is frustration. In some cases, permanent follicle destruction has already occurred before the patient or clinician realizes that the disease is active.

WHAT THE DATA SHOWS

Active scarring alopecia leads at 54.8%. Severe alopecia areata follows at 23.5%, refractory androgenetic alopecia at 15.7%, and female pattern hair loss at 6.0%.

THE INSIGHT

That urgency is consistent with existing specialty guidance. The AAD recommends that patients with active scarring alopecia be referred to a dermatologist for evaluation and treatment. Dermatologists recognize it is also where standard pathways already break down. Clinicians, diagnostic criteria, therapy, disease activity, and clinical differentiation remain unclear.

WHY IT MATTERS

Scarring alopecia is the subgroup where clinicians most often say the pathway is unclear. Respondents are not simply asking for more information. They are asking for a clearer way to decide when a patient is active and when to act.

Subgroup that most changes treatment pathway



07 LONG-TERM OUTCOMES · ADHERENCE · PATIENT EXPECTATIONS

Long-term persistence is supported by multiple signals, not density alone.

51.2% select objective density improvement, but robust shedding, patient-reported benefit, and stable disease activity also matter.

SUMMARY

Hair loss outcomes can be difficult to make clear. Regrowth may be gradual, subjective, and slow to meaningful to assess, and different clinicians demand different endpoints.

WHAT THE DATA SHOWS

Objective density improvement is selected by 51.2%, reduced hair shedding by 39.2%, patient-reported benefit by 35.3%, and stable disease activity by 29.4%. Respondents could choose up to three factors.

THE INSIGHT

Clinician judgment is not a proxy for a patient's opinion. The evidence because a single endpoint can underestimate benefit, particularly in conditions where shedding may be the most visible first improvement.

WHY IT MATTERS

Standardized endpoints, for example, dermatologist-, shedding diaries, and patient-reported outcomes, can help clinicians and trials compare treatments more consistently. They are not a substitute for clinical judgment, but they can help reduce variation in how success is defined.

What most clinicians monitoring long-term hair loss therapy

51.2%	Objective density improvement
39.2%	Reduced hair shedding
35.3%	Patient-reported benefit
29.4%	Stable disease activity

08 AUTHORITATIVE · TOP CLINICIAN RECOMMENDATIONS · RANKING

Unrealistic treatment expectations outweigh affordability, limited diagnostics, and adherence as the main management limitation.

by 41.9% clinicians, treatment expectations as the single biggest limitation.

SUMMARY

Hair loss care is unusually exposed to visual self-monitoring. Patients can notice small changes in shedding, density, or scalp coverage long before a clinician can confirm whether a treatment is working.

WHAT THE DATA SHOWS

Unrealistic treatment expectations lead at 41.9%. Cost / treatment adherence follows at 27.9%, treatment adherence at 23.7%, and limited diagnostics at 6.5%. Respondents could choose up to two factors.

THE INSIGHT

Expectation setting addresses the earlier phenotype result. Scoring diagnosis is not a problem; it is a problem. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. For clinicians, diagnostic criteria, disease activity, and clinical differentiation remain unclear.

WHY IT MATTERS

A premium expectation conversation should not be limited by time issues. For example, a 15-minute visit may not be enough to discuss realistic timelines, likely side effects, and what to do if shedding increases. Clinicians who do not plan for this risk disappointment. This is a practical, implementable solution that leverages your existing workflow.

What most limits effective hair loss management

Unrealistic treatment expectations	41.9%
Cost / treatment adherence	27.9%
Treatment adherence	23.7%
Limited diagnostic resources	6.5%

09 FUTURE · TRICHOSCOPY · AI · TELEMEDICINE

Future optimism is concentrated in new therapies, while AI-assisted trichoscopy ranks far behind.

41.2% prioritize therapeutic innovation; 18% prioritize AI; therapeutic categories show limited overlap.

SUMMARY

The most important opportunity is most likely if the goals should not directly be developed that could most transform hair loss care.

WHAT THE DATA SHOWS

Regenerative therapy leads at 41.2%, future biologic therapies at 29.4%, and new treatments for alopecia areata at 17.6%. AI-assisted trichoscopy is last at 11.8%.

THE INSIGHT

Respondents are primarily looking for better therapeutic options rather than better diagnostic tools. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. For clinicians, diagnostic criteria, disease activity, and clinical differentiation remain unclear.

WHY IT MATTERS

The 18% category means current context. Job pathway therapies are already part of current alopecia areata treatment in the United States, including FDA-approved JAK inhibitors. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. For clinicians, diagnostic criteria, disease activity, and clinical differentiation remain unclear. The low AI score does not mean clinicians reject technology. It may mean they do not see AI as the most urgent unmet need.

Development that could most transform hair loss care within 5–10 years

AREAS REMAINING LONG

The difficult cases are difficult because more than one problem is often present at the same time.

The open-ended question asked clinicians to describe a challenging hair-loss case, which they had seen recently, and what made it complex and unresolved. Responses were grouped into 6 categories. Each category is described below.

Refractory androgenetic alopecia
Cases where standard treatment, such as topical minoxidil, finasteride, and oral minoxidil, have failed to produce adequate regrowth.

Scarring disease and irreversible loss
Cases in which inflammation has already caused permanent follicle destruction, and the clinician's main goal is to prevent further loss.

Overlapping diagnoses
Cases where androgenetic alpecia and scarring alopecia, or alopecia areata patterns, appear together, making it difficult to assign a single diagnosis.

Diagnostic uncertainty
Presentations, whether clinical, trichoscopic, or biopsy findings, that were unclear even after the clinician had completed their usual workup.

Patient preferences and adherence
Cases where the clinical plan was reasonable, but the patient declined, delayed, or could not follow through with treatment.

Prior treatment complexity
Patients who had already been seen by other clinicians and had received multiple treatments before arriving at the current clinician's practice.

What the open responses add: The quantitative survey shows which clinicians usually see and what they usually do next. The qualitative comments show why those decisions are difficult. The most frequent theme was that clinicians are unsure of the diagnosis, unaffected by pain, and responding when the patient is willing and able to continue.

PEOPLE ALSO ASK · TOPIC FAQ

Frequently asked questions about hair-loss diagnosis and treatment.

What is trichoscopy, and how is it used in hair loss diagnosis?
Trichoscopy is a magnified examination of the scalp and hair, usually with a dermatoscope or smartphone attachment. It helps clinicians look for patterns that may not be visible on routine examination, such as changes in hair shaft thickness, scaling, inflammation, or signs of scarring. It can help distinguish between androgenetic alopecia, telogen effluvium, alopecia areata, and scarring alopecia.

When is a scalp biopsy needed for hair loss?
A scalp biopsy is usually considered when the diagnosis is unclear, when scarring alopecia is suspected, or when clinical and trichoscopic findings do not match. It is most useful when it can change treatment, such as when inflammation is present and active treatment may be needed to prevent permanent follicle loss.

Is topical minoxidil a common treatment for early androgenetic alopecia?
Yes. Topical minoxidil is widely used as a first-line treatment for androgenetic alopecia. Clinicians and patients must consider that it can take several months before any benefit is seen, and that stopping treatment can lead to shedding of newly regrown hair.

How long does it take to see results from hair loss treatment?
For patient hair loss, either improvement may take several months. American Academy of Dermatology patient guidance notes that patients should set clear goals and be aware that early shedding can sometimes occur during the first few months of treatment.

How are scarring and non-scarring alopecia differentiated?
Scarring alopecia is characterized by permanent loss of follicles and visible scarring on examination. Scarring, or chemical, alopecia follicles may appear normal, while non-scarring alopecia may show preserved follicular openings and no scarring.

Are JAK inhibitors now used for alopecia areata?
Yes. Oral JAK inhibitors are now approved for alopecia areata in the United States, and JAK pathway therapies have received FDA approval for severe alopecia areata in adults, and several agents are under development. Treatment choice depends on disease severity, age, comorbidities, and local regulatory status.

DIAGNOSTIC PRACTICE GAPS

What this means at four points in the hair-loss care pathway.

The survey suggests the problem is not a lack of diagnostic or scarring alopecia between diagnostic confidence, treatment adequacy, disease urgency, and patient engagement rather than the missing work on essential evidence.

At the first evaluation
Androgenetic alpecia is the default. The most common entry point is also the most dangerous assumption. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. A biopsy could change management.

When hair loss persists
Escalation is the norm. Most clinicians add therapy or escalate their medical treatment when hair loss persists. Only a minority would move to biopsy diagnosis. That is clinically important because "not improving" can mean several different things, including poor adherence, misdiagnosis, or a scarring process that is progressing.

When starting treatment
Expectation setting is the biggest gap. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who do not plan for this risk disappointment.

During long-term follow-up
Success is more than density. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited.

CLOSING PERSPECTIVE

The next improvement in hair-loss care may come from making the pathway more explicit, not simply making the treatment stronger.

This survey suggests that clinical judgment is not the problem. Most clinicians can identify androgenetic alopecia, recognize when trichoscopy or biopsy is appropriate, and start a reasonable first treatment. The harder problem is what happens next. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited.

That is why the most practical improvements may be simple. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited.

The data also show that patient engagement is the central challenge in long-term care. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited.

Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited.

Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited.

Help future dermatology insights reflect the cases that do not fit a simple hair-loss pathway.
Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited. Clinicians who are not interested in trichoscopy or biopsy often overlook treatment adherence limited.

Join the Panel

REFERENCES AND DATA SOURCE

1. Alkhalifah A, Alsantali A, Wang E, McElwee KJ, Shapiro J. Classification of alopecias in patients with alopecia areata. Arch Dermatol. 2010;146(7):749–757.
2. Price VH. Treatment of hair loss. New England Journal of Medicine. 1999;341(13):964–973.
3. Trüeb RM. Pharmacological and nutritional treatment of female androgenetic alopecia. Int J Trichology. 2012;4(3):147–154.
4. Olsen EA, Hordinsky MK, Whiting DA, et al. The importance of dual-energy X-ray absorptiometry results in female pattern hair loss. J Am Acad Dermatol. 2002;46(1):37–47.
5. Rossi A, Cantisani C, Melis M, Iorio A, Scarnò M, Calvieri S. Trichoscopy of androgenetic alopecia: a review. Dermatology. 2012;224(1):1–7.
6. Vañó-Galván S, Pirmez R, Hermosa-Gelbard Á, et al. Androgenetic alopecia in women: a review. J Eur Acad Dermatol Venereol. 2020;34(1):3–14.
7. Lee WS, Lee HJ, Choi GS, Ro BI, Kang H, Kim BJ, Kim MN. Clinical characteristics of diffuse alopecia: clinical and trichoscopic findings. J Eur Acad Dermatol Venereol. 2013;27(6):e274–e281.
8. Trüeb RM, Rezende HD, Dias MFRG, Kasprzak M. Oral minoxidil for androgenetic alopecia in men: a review. Int J Trichology. 2021;13(3):93–101.
9. Trüeb RM, Rezende HD, Dias MFRG, Kasprzak M. Oral minoxidil for androgenetic alopecia in women: a review. Int J Trichology. 2021;13(3):102–110.
10. Trüeb RM, Rezende HD, Dias MFRG, Kasprzak M. Oral minoxidil for androgenetic alopecia in children and adolescents: a review. Int J Trichology. 2021;13(3):111–118.

© 2026 Medicus LLC. All rights reserved. Unauthorized reproduction, distribution, or commercial use of this report is prohibited. Permitted uses include personal research and citation with attribution. Medicus LLC is not responsible for decisions made based on this report. Survey results are subject to sampling and response bias. Terms of Use · Privacy Policy · Cookie Policy