Migraine Treatment and Stress: Why Better Control Needs More Than Pain Relief

Audience-centered study of acute stress, acute treatment, medication overuse, prevention, access, and everyday functioning shape the migraine care journey

STEVE HOFFER · KEVIN HELM · CHRISTEN SAP · PRESENTER · KEVIN · STEVE HOFFER

148 Participants

6 Questions

Questions represented over 100 Likert, Yes/No, and Other / Write-in responses

78.9% Reported stress as a migraine trigger

5618562 Migraine days tracked

BEAT THIS REPORT

Migraine control is not only about how much the head hurts

For many people, migraine management sits at the intersection of pain, uncertainty, stress, work, treatment access, and the ability to live a normal life. This report looks at how patients describe their migraine experience, where their current care falls short, and what they believe would help them feel more in control. The findings point to a clear conclusion: effective migraine care must address not only pain relief, but also the factors that make attacks more frequent, more disruptive, and harder to manage.

This report summarizes these patient voices through the experiences reported by migraine patients across six countries. It follows the patient journey from what sustains attacks to what prevents them, and it highlights the gaps between what patients need and what current care models provide.

METHODOLOGY

How to read the patient data

The survey includes 148 records, of which 105 were complete for a 70.9% completion rate. Respondents were represented across the USA, Canada, Mexico, Brazil, Argentina, and Chile. Participants reported tracking migraine days in their personal calendars before completing the survey. Multiple-choice questions were presented in the source data and are reflected in the percentages used throughout this report.

The survey response export is the numerical record of both. Demographics fields, required-text checks, and honeypot fields are excluded from the analysis. Open-ended responses are used as qualitative context only and are not converted into percentages.

EXECUTIVE SUMMARY

Six signals that define the migraine care gap

01
Some ate inside the migraine cycle
53.8% say stress or anxiety most often worsens migraine, and 38.4% say it makes scheduled work harder to complete.

02
Everyday headaches trigger action
69.2% say worsening of ordinary headaches is a sign migraine is getting worse.

03
Acute care starts with what is accessible
49.6% use over-the-counter pain medicine first during an attack.

04
Poor control does not always lead to escalation
44.7% say the medicine does not escalated after ineffective relief.

05
Function defines treatment value
44.6% say better working hours themselves staying on preventive treatment.

06
Patients want more personalized prevention
36.3% choose more personalized preventive treatment as the future improvement that would help most.

PATIENT JOURNEY STAGE 1

A migraine diagnosis does not always translate into active treatment

26.8% of respondents describing their current migraine experience say they are diagnosed but currently untreated.

CONTEXT

Migraine can be the beginning of care, but it does not guarantee that the person has a treatment plan they are using. Some patients may be between treatment options, may not start medication, may have access barriers, or may be managing symptoms on their own.

WHAT THIS DATA SHOWS

31.4% are diagnosed and currently receiving treatment, 31.7% are recently diagnosed or starting treatment, and 26.8% are diagnosed but currently untreated.

THE INSIGHT

Diagnosis is not a linear progression of possible active treatment stages it represents. The survey answer options say, but the findings show that an appointment booked, treatment was effective, and lack of escalation show several forms of choice that are undiagnosed.

WHY IT MATTERS

Any migraine care improvement pathway needs an answer to a simple, foundational question: Is the patient receiving care, or are they waiting for a option? That may involve acute treatment, prevention, non-drug strategies, mental health support, or referral depending on the individual situation.

STAGE 2 · TRIGGERS

Stress is the factor patients most connect with worsening migraine

Tied to sensory stress or anxiety, and stream of sleep disruptions, hormonal changes, and women's triggers.

CONTEXT

Migraine triggers are individual. Stress, sleep disruption, hormonal changes, weather, food, and sensory overload can all appear in patient descriptions. Current American Migraine Foundation guidance maintains that triggers may influence attacks but are not the underlying cause of migraine.

WHAT THIS DATA SHOWS

53.8% choose stress or anxiety as the factor that most often worsens migraine, followed by sensory stress at 15.4%, hormonal changes at 13.0%, and light, noise, or other triggers at 9.3%.

THE INSIGHT

Stress is not merely a background lifestyle issue in this dataset. It is the most commonly identified trigger, and it sits at the center of a cycle that many patients describe: stress causes attacks, attacks create more stress, and the cycle can be hard to break without treatment that addresses both.

WHY IT MATTERS

Patients benefit from clear guidance on what changes should trigger a call or visit to their clinician. Without that guidance, they may wait too long or escalate on their own pattern. Better control therefore, is rapidly recovering symptoms should not be simply be assumed to be routine migraine.

STAGE 3 · ACUTE TREATMENT

A change in the headache pattern is the strongest prompt for further evaluation

69.2% say worsening or changing headaches prompted further evaluation.

CONTEXT

People with migraine can develop migraines often learn what their usual attacks feel like. A change in the familiar pattern can create uncertainty and may push someone toward a new diagnosis or treatment.

WHAT THIS DATA SHOWS

Among those who sought treatment beyond 30 OTC, increasing severe-related disability is selected by 23.0%, new neurological symptoms by 11.7%, and 10.9% say no further escalation was needed.

THE INSIGHT

Patients appear to experience most strongly when migraine stops behaving like their usual migraine. The second-largest signal is also new to the person, not necessarily disability. That means change can be clinical, functional, or both.

WHY IT MATTERS

Patients benefit from clear guidance on what changes should trigger a call or visit to their clinician. Without that guidance, they may wait too long or escalate on their own pattern. Better control therefore, is rapidly recovering symptoms should not be simply be assumed to be routine migraine.

STAGE 4 · ACUTE TREATMENT

During an attack, the most common first step is over-the-counter pain medicine

49.6% use over-the-counter pain medicine first, compared with 36.2% who use a triptan first.

CONTEXT

Acute migraine treatment is important for what a person has been prescribed, when they take it, and how well it works for their specific attack. Timing may explain treatment can be taken.

WHAT THIS DATA SHOWS

49.6% say they use over-the-counter first. Triptans are selected by 36.2%, gepant or CGRP-targeted treatments by 7.2%, and ditans by 1.4%, or neither by 5.8%.

THE INSIGHT

Pain relief is the most accessible first response, but it is not always the most effective. The higher-tier treatments are available, but they are not the default for most patients in this dataset.

WHY IT MATTERS

A migraine plan should make acute treatment instructions clear before the next attack arrives, what to take, when to take it, what to avoid combining, and when frequent acute treatment use should trigger a review of preventive care.

STAGE 5 · MEDICATION OVERUSE

Stress remains uncontrolled even after treatment enters the picture

35.4% directly uncontrolled stress or anxiety as the single biggest limitation of current migraine treatment.

CONTEXT

Treatment limitations can come from inconsistent use, incomplete symptom relief, tolerability, or access. Medication overuse headache, a common medication overuse, is a serious medication-related risk that can worsen migraine care.

WHAT THIS DATA SHOWS

Stress or anxiety remains uncontrolled for 35.4%, incomplete symptom relief for 34.3%, medication side effects for 15.3%, and medication overuse or dependence for 5.6%.

THE INSIGHT

Stress is the factor with the largest symptoms and the treatment limitation selected. The pattern suggests that many patients can manage the physical and neurological parts of the migraine experience remain intertwined even after treatment begins.

WHY IT MATTERS

The Overcome Migraine Foundation describes migraine and mental health as closely connected and notes that anxiety, stress, unpredictability, and disruption can all affect migraine. That helps explain why patients may feel that migraine is psychological, or that treatment works for the pain but not the stress that triggers it. A prevention plan that addresses only one of these two problems is unlikely to produce a major leap for patient experience improvement.

PATIENT JOURNEY STAGE 6

When migraine control is inadequate, the next step is not always escalation

49.7% say treatment was not escalated after inadequate migraine control.

CONTEXT

Treatment escalation is intended to reduce attack frequency, severity, or disability over time. Current American Headache Society resources emphasize matching prevention to migraine burden, patient preferences, medical history, and treatment goals, not escalation alone.

WHAT THIS DATA SHOWS

Of 105 reports relating to an unanswered, 13.0% selecting CGRP-targeted therapy, and 5.8% selecting botulinum toxin. The largest group, 49.7%, did not escalate after inadequate control.

THE INSIGHT

Not all patients who are offered any treatment was not escalated. The reasons may range from clinical appropriateness to cost, access, tolerability, or difficulty, access, or the timing of follow-up. What it does show is a subtle but important gap between what patients are told to do and what they actually do. Inadequate control can become the new normal rather than a signal to act.

WHY IT MATTERS

Patients should have clear treatment success is supposed to look like and when to call their clinician if it is not working. Without that benchmark, they can underestimate how much worse they have become or how much help is available.

EXECUTIVE SUMMARY

Anxiety, sleep, low mood, and stress can all complicate migraine care

51.9% select stress-related symptoms and 43.5% anxiety or panic symptoms as the coexisting issue that most complicates care.

CONTEXT

Migraine is often experienced alongside sleep problems, anxiety, depression, and stress. These conditions can affect how symptoms are perceived, how prevention plans fits them, and how easy it is to follow a treatment plan.

WHAT THIS DATA SHOWS

Stress-related symptoms screening leads, burnout, at 51.9%, followed by anxiety or panic symptoms at 43.5%, depression or mental at 31.5%, and dependence on the result at 31.5%.

THE INSIGHT

The migraine experience is often overloaded by changes in migraine days, but symptoms that extend the disease through missed, disrupted days, fatigue, anxiety to light or sound, recovery time, and anxiety about the next attack.

WHY IT MATTERS

An OTC-only strategy can be tempting as the main influence in staying with prevention treatment. Faster monthly migraine days follow a 20.5%, more than 3 times the 6.4% who report 15 to 20 days. Patients who are not yet at that threshold may not see the urgency.

PATIENT JOURNEY STAGE 7

Patients judge preventive treatment by whether life becomes easier

44.6% say better daily functioning is the biggest influence on staying with preventive treatment.

CONTEXT

Preventive migraine treatment is often evaluated by changes in migraine days, but patients may experience benefits in other ways. Improved sleep, reduced fatigue, anxiety to light or sound, recovery time, and anxiety about the next attack can all matter.

WHAT THIS DATA SHOWS

44.6% choose better daily functioning as the main influence on staying with preventive treatment. Fewer monthly migraine days follow at 23.2%, more than 3 times the 6.4% who report 15 to 20 days.

THE INSIGHT

The most effective preventive appear to functional, not symptomatic. Patient appear to judge treatment success by how much it gives back to their life. Fewer days matter, but only if the person can actually work, socialize, and enjoy their time without constantly worrying about the next attack.

WHY IT MATTERS

If a patient is having fewer migraine days but still cannot work, sleep, exercise, or plan their life around the possibility of an attack, they may not feel that their treatment is working. Clinicians should therefore ask not only how many attacks occurred, but what the person can do now that they could not do before.

STAGE 8 · FUTURE CARE

The next migraine improvement patients want is more personalized prevention

35.0% choose personalized preventive treatment, while 34.3% say cost or insurance coverage is the biggest current access barrier.

CONTEXT

Personalized options is why varied in a patient care team it affects to fit into the migraine care and treatment needs. Across formats can help migraine care and support for sustaining mental health needs.

WHAT THIS DATA SHOWS

Cost or insurance coverage is the leading access barrier at 34.3%, followed by time or appointment burden at 24.7%, lack of information at 21.5%, and limited options availability at 19.4%. Among those who sought treatment beyond OTC, 35.0% choose personalized preventive treatment as the future improvement that would help most, and 23.0% choose faster access to newer migraine therapies.

THE INSIGHT

Patients are not asking for novelty alone. The leading priorities are personalization and integration. Faster access to newer therapies matters, but it ranks below having prevention tailored to the person and having migraine care connected with mental health support.

WHY IT MATTERS

Personalization can only be judged by how far how many options and not need to another the right option can be matched to the right person without unnecessary cost, delay, or fragmentation.

MULTIVARIATE ANALYSIS

When stress rises, patients look for quiet, control, and room to recover

Respondents were asked to describe a time when anxiety or stress worsened migraine and what would have helped. The comments do not represent a controlled quantitative sample, but they add practical detail to the survey numbers.

Quiet and sensory reduction
Several people described needing a dark, quiet room, minimal lights, less noise, and fewer sensory demands during an attack.

Sleep and exercise that
Some linked stress with disrupted sleep and said regular rest, routine, or exercise could help prevent attacks from building.

Environmental control
Others described needing to leave, step away, or change their environment when migraine symptoms became overwhelming.

Coping tools
Breathing exercises, meditation, understanding coping mechanisms, and knowing what to do were also described as helpful.

Treatment that prevents the attack
Some respondents wanted a more effective treatment option or preventive medication when they could feel early building toward a migraine.

Work pressure
Deadlines, meetings, noise, crowded rooms, and high-pressure work environments were repeatedly described as times when migraine was harder to manage.

Different experience with loss
A few respondents described migraine during grief or anxiety contexts. One noted that the situation may have been more about anxiety and grief, and not a migraine at all.

The qualitative comments reinforce a central point from the quantitative data: patients often experience migraine management as a need to control both the attack and the conditions around the attack.

PATIENT JOURNEY LENS

What a more usable migraine care pathway could look like

The survey suggests four places where patients could be better supported if care were more responsive and better coordinated.

1. Before the next attack
Patients need clear guidance on when worsening is not to be early, what they should do when they feel a change coming, and when to call their clinician.

2. When control is still inadequate
Know the red flags. The red flags should define what prevention, treatment adjustment, or referral options are available.

3. When the patient changes
Changes in headache pattern, new neurological symptoms, new symptoms, or increasing disability should lead to a more comprehensive plan.

4. When life is still not working
Measure what matters. Daily function, stress, sleep, pain, work, and social participation should be part of every migraine visit, not just migraine days.

CLOSING PERSPECTIVE

Better migraine care is not one more treatment. It is a clearer path through the whole experience

This survey shows migraine as a condition that moves through several dynamic parts at once. An attack itself. But factors that make attacks harder, the uncertainty of what to do next, and the ways migraine can interfere with work, relationships, and the practical difficulty of accessing care.

The central tension is that stress appears both at the most important place. It is the leading factor patients say worsens migraine, and it is the leading reason they say remains uncontrolled during treatment. That does not make migraine a stress disorder. It does suggest that for many patients, the two are inseparable in practice.

The treatment pathway also shows room for improvement. Most patients who seek care beyond OTC medication do not receive a triptan on the first attempt. Many people who try a preventive do not stay on it long enough to see whether it can help them. And a substantial group of patients who are not receiving treatment at all may be underestimating how much migraine is affecting their lives because they have normalized it. Together, these findings point to a system that is not always failing patients, but is often not responsive enough to their changing needs.

Perhaps the most valuable takeaway from this survey is that patients want more than relief. They want control. They want to know when to act, when to wait, when to escalate, and when their plan is working. They want to feel that they have a plan for how to live with migraine, not just how to survive the next attack.

Make treatment goals explicit
Define what success looks like for each patient: fewer attacks, better function, less anxiety, and sustainability.

Build in a review point
Schedule a check-in when a new treatment starts so that patients know when to call their clinician if it is not working.

Treat stress as part of the plan
Stress management, sleep, and coping strategies should be integrated into preventive migraine treatment.

Bring stress and sleep into the conversation
Ask about stress, sleep, and daily functioning at every visit, not only when migraine days are high.

Share what migraine care feels like in real life

The Chronicity Project collects patient stories to help clinicians understand what prevention, rescue, support, and day-to-day care.

Join the Project

PEOPLE ALSO ASK

Frequently asked questions about migraine treatment, stress, and prevention

Can stress trigger a migraine attack?
Stress can be a trigger for some people with migraine, although triggers differ from person to person. Stress is not considered the underlying cause of migraine. Treating stress may help reduce attack frequency for patients who identify it as a trigger.

What should I do if my migraine treatment is not working well enough?
Keep a migraine diary and bring it to your next appointment. Migraine management can include reviewing the diagnosis, how acute medicines are being used, and whether preventive treatment may be helpful. Medication overuse can also make headaches more frequent.

When is preventive migraine treatment considered?
Preventive treatment may be considered when migraine attacks are frequent, disabling, prolonged, or difficult to manage with acute treatment alone. The right option depends on your migraine pattern, other health conditions, preferences, and side effects.

Can anxiety or poor sleep make migraine harder to control?
They can be important parts of the migraine picture. Stress, anxiety, and disrupted sleep may interact with migraine symptoms and daily functioning. A care plan may need to address migraine attacks and the factors that make them harder to live with.

What is medication overuse or rebound headache?
Medication overuse headache can develop when pain medicines are used too frequently for too long. It can feel like yourself recently acute treatment you often, discuss it with a clinician. Reducing medication use may require a structured plan, not sudden stopping.

How can I prepare for a migraine appointment?
A simple record of migraine days, severity, disability, medicines used, side effects, possible triggers, and how attacks affect work or daily life can help your clinician understand your pattern. Bring this information to your next visit.

References and external content

1. American Migraine Foundation. Migraine Triggers. https://americanmigrainefoundation.org/resource-library/migraine-triggers/
2. American Migraine Foundation. Migraine Prevention. https://americanmigrainefoundation.org/resource-library/migraine-prevention/
3. American Headache Society. Pharmacologic Treatment for Migraine Prevention in Adults Practice Guideline Recommendations. Published August 15, 2025. https://americanheadachesociety.org/news/pharmacologic-treatment-for-migraine-prevention-in-adults-practice-guideline-recommendations/

External sources provide context; MRM Chronicity findings are interpretations unless directly from patient-reported responses.

© 2026 MRM Group
All Rights Reserved. Chronicity
Privacy Policy · Terms of Use · Contact