Surgical Site Infection Prevention: Where Patient Risk Meets Protocol Adherence

THE GLOBAL GUIDE TO INFECTION CONTROL · EXPERT CONSENSUS REPORT

Surgical site infection prevention fails, whenever timing, wound classification, bundle adherence, ostomy control, and surgical team coordination don't work together.

INFOGRAPHIC · PREVENTION STRATEGIES · PREVENTION SUCCESS · SURVEY FOR 50 & SHARPE CONTROL

202	6	71.3%	900180
Surgical Procedures	Countries Represented	Of Control Risk	Of Control Risk

Survey Points: 202 survey records with 6 LTIs completed rate. Country: Indiana Department for 2021, UK, Canada, Italy, France, and Germany. Conference Summit Center map from 2021 to the Guarantee. Internal Improvement, Improvement, Improvement, and Surgical Safety and Control Risk.

WHY THIS MATTERS

SSI prevention becomes difficult when standardized safeguards meet patients and procedures that are not.

As in surgery, infection prevention is a daily preventive discipline. Surgical site infection prevention and classification, models through prophylaxis, skin preparation and operative technique, and can rapidly become a diagnostic and source control problem in surgical patients.

The WOFAS core survey captures where practice gaps the greatest weight across the pathway. The responses highlight poorly controlled diabetes, contaminated wound classification, and delayed antibiotic prophylaxis as the most frequent risk factors, while the overarching challenge of making preventive practices consistent and reliable in real-world surgical settings.

CLINICAL CONTEXT

Modern SSI prevention is a bundle of evidence-based steps, but implementation still depends on the operation, the patient, and the reliability of the system around them.

The CDC SSI prevention guideline remains a core UK reference, while the SHEA/IDSA/APIC post-care update emphasizes evidence-based practices for surgical patients, including preoperative optimization, bundle adherence, and operating room management.

The WHO global SSI guidelines and NICE guidance similarly frame prevention as a system issue, one , and postoperative periods. None of these recommendations can be applied in isolation. Surgical wound management, hand hygiene, skin preparation, and wound closure must change the practical pathway.

The WOFAS survey reflects this gap between recommendation and execution. It does not test which intervention is clinically superior. It shows which risks and decisions surgeons in this respondent group say most influence their practice and the patient.

EXECUTIVE SUMMARY

Six findings that show where SSI prevention shifts from protocol to risk-sensitive surgical judgment.

01

69.7%

Protocol adherence is the dominant countermeasure gap

Protocol non-adherence is the most selected limitation to delivering consistent SSI prevention.

02

52.9%

Poorly controlled diabetes is the patient profile most likely to intensify SSI prevention

03

51.9%

Poorly controlled diabetes drives prevention intensity

Diabetes is the patient profile most often selected as increasing SSI prevention effort.

04

43.8%

Contaminated or dirty surgery changes strategy most

Wound contamination leads the operational scenarios that most alter SSI prevention.

05

49.4%

Timing is the main evaluation of surgerical drug SSI

Timing is the most selected real-time step in a drug's surgical site infection is suspected.

06

34.2%

Personalized protocols lead the future outlook

Personalized surgical protocols is the development most often expected to transform SSI prevention.

SECTION 01 · RISK FACTORS

Poorly controlled diabetes is the patient profile most likely to intensify SSI prevention

51.9% say poorly controlled diabetes most increases their SSI prevention efforts.

SUMMARY

Poorly controlled diabetes leads by from a different baseline when wound healing, immune response, or metabolic control is already compromised. Diabetes is a well-established risk factor for surgical site infection, and the survey results show that clinicians adjust their prevention practices accordingly.

WHAT THE DATA SHOWS

Poorly controlled diabetes leads at 51.9%, followed by immunosuppression at 43.1%, current obesity at 37.4%, and previous SSI at 31.6%. A separate question on patient factors that increase SSI risk most also selected diabetes at 51.9%.

THE INSIGHT

The survey shows that diabetes is not just a risk factor, but a pattern of risk that clinicians count as a prompt to change their practice. The consistency between the two questions suggests that diabetes is a practical trigger for more intensive SSI prevention, not just a theoretical concern.

WHY IT MATTERS

For surgical teams, this emphasizes to not the one metabolic case determines whether surgery proceeds. It is that poorly controlled diabetes can begin a cascade of additional precautions, from timing of antibiotics to wound dressing choices to the overall intensity of the perioperative bundle.

Patient profiles that most increase SSI prevention efforts



SECTION 02 · PREVENTION STRATEGIES

Contaminated wound classification changes the prevention plan before surgery starts

43.8% say contaminated wound classification most changes their preoperative SSI strategy.

SUMMARY

Preoperative SSI planning is rarely about anticipating recorded number. It often becomes primarily an operative choosing between bundle to the wound classification. Contaminated or dirty surgery changes the threshold for antibiotics, skin preparation, and wound management, and the survey results show that clinicians adjust their plans before the same room safety practices are used.

WHAT THE DATA SHOWS

Contaminated wound classification leads at 43.8%, followed by open ground to wound at 41.2%, MRSA colonization at 37.7%, and long operative duration at 34.2%.

THE INSIGHT

The ranking shows that surgeons weight most operative contamination slightly more than patient-level factors when they plan SSI prevention. This suggests that preoperative strategy is dominated not only by who the patient is, but by how contaminated the surgical field is expected to be.

WHY IT MATTERS

For surgical teams, this emphasizes the importance of accurate, consistent documentation of wound classification. It also shows why perioperative preparation for contaminated cases must be more than a checklist. Antibiotic timing, skin preparation, and wound management all depend on how the surgeon and team categorize the risk before the first incision.

Findings that most change preoperative SSI strategy



SECTION 03 · PRE-INCISION PREVENTION

Timely antibiotic prophylaxis remains the pre-incision measure surgeons prioritize most

52.9% prioritize timely antibiotic prophylaxis before incision.

SUMMARY

The results below reinforce complete standard prevention tasks as a short preoperative window. Timely antibiotic prophylaxis, skin preparation, antibiotic redosing, equipment readiness, and team checks all compete for attention, which means reliability can depend as knowing the recommendation.

WHAT THE DATA SHOWS

Timely antibiotic prophylaxis leads at 52.9%, Chlorhexidine alcohol skin preparation at 48.5%, and Appropriate antibiotic redosing at 41.1%. Equipment readiness checks is at 1.2%.

THE INSIGHT

The ranking suggests that many surgeons prioritize defining anatomy and timing over the preparation of the surgical field. Skin preparation is also a key step, but it is often treated as a routine task rather than a decision point. Redosing is a visible, access to the room, and the likelihood that longer OR intraop management.

WHY IT MATTERS

Even SSI management is often a control system problem as much as an individual decision. Timely antibiotic prophylaxis, skin preparation, and antibiotic redosing are likely to be multifaced and whether changes in operative management should have made to the planning.

Pre-incision prevention measures prioritized by surgeons



SECTION 04 · INFECTION CONTROL

SSI prevention failure is seen as a combination of uncontrolled patient risk and process breakdown

42.6% identify uncontrolled patient risk as the most common driver of SSI prevention failure.

SUMMARY

Even with a 202-survey bundle prevention efforts, the breakdown is viewed for a single break may. It shows. Infection risk emerge from the factors, contamination, operative constraints, and the reliability of the prevention practices that are meant to control them.

WHAT THE DATA SHOWS

Uncontrolled patient risk leads at 42.6%, followed by protocol non-adherence at 31.7%, contamination at 12.3%, and resource constraints at 9.7%.

THE INSIGHT

Surgeons do not believe failure only to technical breaches. The largest share of responses points to patient risk that is not adequately controlled before the procedure. The pattern suggests that preventable SSI is often understood as a failure to recognize and manage risk before the operation begins.

WHY IT MATTERS

A useful SSI review needs to separate what was difficult to change from what was simply not done well. Patient optimization, glycemic control, and infection prevention bundle are usually considered before surgery, during operation, and after the procedure bundle was actually completed.

What most commonly drives SSI prevention failure



SECTION 05 · OPERATIVE FACTORS

When deep SSI is suspected, imaging is the most selected first move

49.4% select CT or MRI evaluation as their most common step for suspected deep SSI.

SUMMARY

Deep SSI is harder to diagnose than a superficial infection. Fever, pain, and drainage may not be obvious. Most clinicians in this survey would not wait for a clear clinical picture before ordering imaging. They would start with CT or MRI to look for deep collection, abscess, or hardware involvement.

WHAT THE DATA SHOWS

CT or MRI leads at 49.4%, followed by Wound culture at 28.2%, Empiric IV antibiotics at 16.7%, and surgical exploration at 7.6%.

THE INSIGHT

The ranking suggests that many surgeons prioritize defining anatomy and timing over the preparation of the surgical field. Skin preparation is also a key step, but it is often treated as a routine task rather than a decision point. Redosing is a visible, access to the room, and the likelihood that longer OR intraop management.

WHY IT MATTERS

Even SSI management is often a control system problem as much as an individual decision. Timely antibiotic prophylaxis, skin preparation, and antibiotic redosing are likely to be multifaced and whether changes in operative management should have made to the planning.

Primary next step for suspected deep SSI

SECTION 06 · OPERATIVE FACTORS

Contaminated or dirty surgery and prosthetic implants create the biggest strategic shifts

43.8% say contaminated or dirty surgery most changes their SSI strategy. 38.6% select prosthetic implant surgery.

SUMMARY

Contamination can change the consequences of infection. Contaminated fields may require more aggressive irrigation, longer antibiotic coverage, or earlier source control. Implants can add a new set of considerations complications risk, removal, or prosthetic infection, revision burden, or long-term function.

WHAT THE DATA SHOWS

Contaminated or dirty surgery leads at 43.8%, followed by prosthetic implant surgery at 38.6%, long operative duration at 34.2%, and previous SSI at 31.6%.

THE INSIGHT

The only different mechanisms dominate the responses. Higher contamination at the time of surgery and the presence of foreign material that raises the consequences of infection. Both can change the threshold for more aggressive treatment and specific planning.

WHY IT MATTERS

For surgeons working with infectious disease or stewardship teams, the practical question is not simply whether a case is positive. It is whether the wound, implant, and patient factors together justify a more intensive approach. This can influence the surgical technique and the operative and duration and the team's decision.

Surgical scenarios that most change SSI strategy

SECTION 07 · ANTIBIOTIC MANAGEMENT

Clinical improvement and source control carry more weight than a negative culture alone

35.4% say clinical improvement most supports narrowing or stopping SSI antibiotics. 33.8% select adequate source control.

SUMMARY

Even an SSI is being treated, the decision to narrow or stop therapy is rarely based on one laboratory result. Culture negative, clinicians may continue antibiotics if the patient is still septic, the source has not been controlled, or the wound has a significant infection.

WHAT THE DATA SHOWS

Clinical improvement leads at 35.4%, closely followed by adequate source control at 33.8%. Negative blood culture ranked 3rd at 22.3%, while Persistent fever/sepsis ranked 4th at 8.5%.

THE INSIGHT

The ranking suggests that many surgeons prioritize defining anatomy and timing over the preparation of the surgical field. Skin preparation is also a key step, but it is often treated as a routine task rather than a decision point. Redosing is a visible, access to the room, and the likelihood that longer OR intraop management.

WHY IT MATTERS

Even SSI management is often a control system problem as much as an individual decision. Timely antibiotic prophylaxis, skin preparation, and antibiotic redosing are likely to be multifaced and whether changes in operative management should have made to the planning.

What most supports narrowing or stopping SSI antibiotics

SECTION 08 · CONSISTENCY GAP

Protocol non-adherence is the dominant barrier to consistent SSI prevention

69.7% identify protocol non-adherence as the biggest limitation to consistent SSI prevention.

SUMMARY

Even with a 202-survey written practice and still above, surrounding prevention. Surgical teams, anesthesia, wound care, nurses, and, and may be variable in how consistently they implement the recommendations. Protocol non-adherence is the most selected limitation to delivering consistent SSI prevention.

WHAT THE DATA SHOWS

Protocol non-adherence leads at 69.7%, followed by Resource constraints at 9.7%, and Unfamiliarity with guidelines at 8.5%. Equipment unavailability is at 1.2%.

THE INSIGHT

The gap is striking because resources and time limitations are not yet dominant explanations. Most respondents point to the failure to apply what they already know. This suggests that the real challenge is not awareness, but how to consistently perform the prevention bundle.

WHY IT MATTERS

For surgical teams, this is a practical reminder that SSI prevention is not only a knowledge problem. It is also a workflow problem. The survey suggests that the most important gains may come from making preventive steps more routine, verifiable, and difficult to bypass during a busy surgical day.

What most limits consistent SSI prevention

SECTION 09 · FUTURE OUTLOOK

Surgeons expect personalization to matter more than a single new prevention technology

34.2% select personalized surgical protocols as the development most likely to transform SSI prevention within 3–5 years.

SUMMARY

SSI prevention innovation is moving in several directions at once: better risk prediction, more personalized protocols, and new technologies that can help detect surgical site infection earlier. The survey suggests that clinicians are not waiting for a single breakthrough to change practice.

WHAT THE DATA SHOWS

Personalized prevention protocols lead at 34.2%, followed by antimicrobial surgical coatings at 28.7%, and AI-based risk prediction at 22.4%. Wearable monitoring is at 14.7%.

THE INSIGHT

The ranking suggests that many surgeons prioritize defining anatomy and timing over the preparation of the surgical field. Skin preparation is also a key step, but it is often treated as a routine task rather than a decision point. Redosing is a visible, access to the room, and the likelihood that longer OR intraop management.

WHY IT MATTERS

Even SSI management is often a control system problem as much as an individual decision. Timely antibiotic prophylaxis, skin preparation, and antibiotic redosing are likely to be multifaced and whether changes in operative management should have made to the planning.

Developments expected to transform SSI prevention within 3–5 years

SECTION 10 · PERSONALIZED PREVENTION

Challenging SSI cases represents move from prevention into source control, methodology, wound management, and patient adherence.

The open-ended question asked surgeons to describe a challenging SSI case, what, increased infection risk, how management changed, and what ultimately determined the outcome. Responses were coded into the themes below and ranked by how often they appeared.

Unstable net tissues with wounds that provide the hygiene

Prevention was the most common theme, with surgeons describing how preoperative optimization, perioperative antibiotics, and wound management can be undermined by poor tissue quality, contamination, or delayed recognition of infection.

Source control especially changed management

Many cases required more than prevention alone. Surgeons described drainage, debridement, and source control as the turning points in severe infections. The common message was not simply that SSI can be treated, but that it can change the entire plan for the patient and the wound.

Patient adherence will likely as remained unresolved

Several responses described patients who did not follow wound care, antibiotic, or postoperative instructions, which can, in turn, affect the outcome. These cases underline that SSI prevention is not only a technical problem. It also depends on how well the patient can be supported and followed during recovery.

Multidisciplinary and team coordination limits could the culture

Finally, some cases were complex because of delays in diagnosis, unclear responsibility between surgeons, anesthesia, and wound care teams, or poor communication across different settings. Many clinicians described these as system failures as much as clinical ones.

PROFILE BLOCK 1 OF 2 · TYPE 1 DM

Frequently asked questions about surgical site infection prevention.

What is a surgical site infection?

A surgical site infection is an infection that occurs after surgery in the part of the body where the surgery took place. It can affect the skin, tissue, or even deeper structures around the incision.

When should antibiotics be given before surgery?

Antibiotics are usually given within 60 minutes before the surgical incision is made. This timing helps ensure that enough medicine is present in the tissues when the surgery begins.

How does diabetes affect the risk of surgical site infection?

Diabetes can increase the risk of infection because it can affect how well the body heals and fights infection. Keeping blood sugar levels well controlled before and after surgery is very important.

Why is Staphylococcus aureus decolonization used before surgery?

Some patients carry bacteria like Staphylococcus aureus in the nose without symptoms. Decolonization is sometimes used to reduce the chance that these bacteria will cause infection after surgery.

How is a suspected deep surgical site infection evaluated?

If doctors suspect a deep infection, they may order imaging, such as a CT scan or MRI, to look for collections of pus or other signs of infection deeper than the skin.

When can antibiotics be stopped or changed after a surgical site infection?

Antibiotic decisions depend on the clinical course, culture information, source control, patient factors, and how the wound looks. Stopping or changing antibiotics is usually based on clinical improvement and appropriate source control, not on a single test result.

SECTION 11 · PREVENTION BUNDLE

What this means at four points in the surgical pathway.

The survey suggests that SSI prevention is strongest when the team bundle is stable, while patient and procedural factors can still change the risk.

1. Modify risk before incision

Make risk actionable. Diabetes control, infection status, ostomy, previous SSI, and wound classification should be identified before the operation begins.

2. Separate diagnosis from source control

When deep SSI is suspected, early imaging is important, but the real question is whether the infection can be drained, debrided, or controlled. Antibiotics alone may not be enough.

3. Treat the intra-critical preparation steps

Make the bundle routine. Timely antibiotics, skin preparation, wound protection, and team checks should be consistent enough that they do not depend on memory alone.

4. Review the system, not only the complication

Look for patterns. When SSIs occur, ask whether the problem was patient risk, antibiotic timing, wound care, operative duration, ostomy, materials, and follow-up as the learning must be shared across the team.

CLOSING PERSPECTIVE

The prevention opportunity is not choosing between standardization and personalization; it is making them work together.

The WOFAS survey describes a surgical infection pathway in which control prevention measures remain strongly , yet their implementation changes with the patient and the procedure. Poorly controlled diabetes, wound contamination, previous SSI, and prolonged operative duration are the preoperative factors that most frequently change how surgeons plan for infection prevention.

The strongest operational finding is the size of the preventive gap. Many surgeons in this survey identified protocol non-adherence as the biggest limitation to consistent SSI prevention. This suggests that the most important improvements may not come from new knowledge alone, but from making the right steps easier to perform reliably in busy surgical settings.

At the same time, surgeons are not relying on infection prevention to prevent. Source control, timely antibiotics, and wound management are also central to how clinicians think about SSI risk. The future of prevention will likely depend on combining these approaches: personalizing care where patient risk is high, while keeping the core bundle consistent and easy to follow.

The future signal is therefore balanced: personal is dependable bundle factors. Next personalized protocols extend patient risk, procedure type, microbiology, and recovery data to guide prevention more precisely. Antimicrobial surgical coatings and AI risk prediction may help, but only if they are integrated into existing workflows rather than added as separate tools.

Make risk factors visible early

Use diabetes control, wound classification, ostomy, SSI, and operative duration to identify patients who may need a more intensive prevention plan before the day of surgery.

Audit the prevention bundle

Review antibiotic timing, skin preparation, wound protection, and team checks to find where the routine is breaking down in practice.

Personalize without creating unreliable variation

Use clinical and microbiological data to support better decisions, but keep the core steps consistent enough that they can be delivered reliably.

Help Sharpe surgical safety insights reflect the realities of SSI prevention in practice.

Join the WOFAS Network to share your experiences and help improve surgical safety guidance.

Join the Survey

REFERENCES AND DATA SOURCE

1. Methicillin-resistant Staphylococcus aureus (MRSA) in the United States: Epidemiology and Genetic Relationship to Community-Acquired Staphylococcus aureus. Clin. Infect. Dis. 2005; 41:1131–1136. Epub 2005 Sep 12. doi: 10.1086/444459. PMID: 16163626.
2. Antibiotic Use Hospital and Community. Surgical Site Infection Prevention and Treatment (SSIPT). CDC. Infection Control.
3. Surgical Site Infection Prevention: Where Patient Risk Meets Protocol Adherence. Expert Consensus Report. Sharpe Control Risk. 2021.
4. National Institute for Health and Care Excellence. Surgical site infections: prevention and treatment (NICE CG74). NICE recommendations.
5. Surgical Site Infection Prevention: Where Patient Risk Meets Protocol Adherence. Expert Consensus Report. Sharpe Control Risk. 2021.

Sharpe Control Risk is an independent research and consulting organization specializing in infection control, risk management, and clinical safety. Our reports are based on survey data, expert interviews, and analysis of real-world practice. We do not accept commercial sponsorship for our research publications.

This report is for informational purposes only. It does not constitute medical advice, clinical guidance, or a substitute for professional judgment. Readers should consult qualified healthcare professionals before making decisions about patient care.

© 2026 Sharpe Control Risk. All rights reserved. Reproduction or distribution of this report, in whole or in part, without prior written permission is prohibited.